2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day Resident Posters

Resident Poster Resident

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 003 Category: Clinical Vignette

Residency Program: Ascension Macomb-Oakland
Presenter: Shamaiza Waqas
Additional Authors: Sean Dawes, DO MPH, Jorawar Brar DO, Waqas Abid MD ,Amaraja Kanitkar MD MPH

Atrial septal aneurysm leading to Ischemic stroke : A case report and literature review

Background:

Stroke is the second most common cause of death and disability in the world 13.7 million new cases of stroke were noted in 2016, out of which around 87% were secondary to ischemic strokes. Atrial septal aneurysm (ASA) is a localized saccular deformity of the atrial septum, associated with ischemic stroke independently or in association

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 004 Category: Continuous Quality Improvement/Evidence-based Medicine

Residency Program: Ascension Providence
Presenter: Zachary Bassler
Additional Authors: Avery Mendelson MD; Saif Affas MD; Christopher Hakim MD

Navigating Colorectal Cancer Screening: Insights from a Quality Improvement Study

Background: Colorectal cancer ranks as the second leading cause of cancer-related deaths in the United States, disproportionately affecting individuals aged 65-74. While screening recommendations have shown efficacy in reducing mortality, there exists a concerning rise in incidence among adults aged 45-49. This quality improvement study addresses the need for improved screening compliance, particularly among the age group of 45 to 50 to mitigate the impact of colorectal cancer.

Methods: A quality improvement study was conducted targeting average-risk individuals aged 50-75 in an academic internal medicine clinic. Patient outreach and education, facilitated through virtual visits, aimed to improve colorectal cancer screening compliance. Secondary goals included assessing the performance of academic internal medicine practitioners in colon cancer screening, enhancing patient education on screening modalities, and promoting awareness of the benefits of age-appropriate screening.

Results: Of the 1258 identified patients deficient in colorectal cancer screening, 215 were chosen as the initial study population. A total of 36 orders were placed for patients. There was a slight preference for non-invasive screening during the initial outreach phase of the study. Despite challenges in patient outreach, 30.5% of orders

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster #: 005 Category: Clinical Vignette

Residency Program: Ascension Providence
Presenter: Atris Rose
Additional Authors: Atris Rose; Leo Parsons

George Felemegos

Delayed Brain Abscess Formation Due to Streptococcus Pneumoniae Following Craniotomy in an Immunocompetent Young Adult: A Case Report

Bacterial meningitis presents a life-threatening scenario requiring immediate and comprehensive management, as highlighted in the challenging case of a 23-year-old male who arrived at the Emergency Department (ED) unresponsive, exhibiting fever, headache, and seizures. His medical history included a traumatic brain injury from a previous motor vehicle accident, with subsequent frontal craniotomy in Colombia. The patient's symptoms, including a recent tonic-clonic seizure, raised concerns for acute bacterial meningitis and prompted diagnostic and therapeutic interventions to address the intricacies of this condition and its potential intracranial complications.

He was unresponsive upon admission, subsequently intubated and sedated, and transferred to ICU, after a lumbar puncture confirmed bacterial meningitis. Treatment involved Decadron, Ceftriaxone, cefepime, and vancomycin. Preceding the lumbar puncture, a CT head revealed post-op changes from a frontal craniotomy with bilateral encephalomalacia, ruling out hemorrhage. Neurology consultation and EEG findings excluded ongoing seizures, while MRI indicated bifrontal cerebritis with a possible abscess. Neurosurgery, concerned about a potential abscess, added Flagyl to the antibiotic regimen. Blood cultures identified Streptococcus pneumoniae. EVD placement addressed ventriculitis, leading to clinical improvement, successful extubation, and subsequent identification of a CSF leak. ENT consultation, confirmed by a CT cisternogram, diagnosed defects in the cribriform plate and sphenoid bone, prompting surgical intervention with right ethmoidectomy and mucosal grafting. Pathology revealed a necrotizing ulcer, and a follow-up MRI showed resolution of cerebritis. The patient completed a 4-week course of Rocephin, with plans for repeat imaging and a Prevnar 20 vaccine before discharge.

In a retrospective study spanning 1991 to 2005, involving 2111 neurosurgical procedures, the incidence of postoperative central nervous system infection (PCNSI) was less than 1%, irrespective of factors like cerebrospinal fluid leakage and diabetes. Targeted prophylaxis against pathogens, especially Staphylococcus aureus and Propionibacterium acnes, is crucial in high-risk cases. Another study examined 44 patients with cerebral abscesses post-brain surgery, emphasizing their rare and severe nature. Tumor resection (61%) was common, with Staphylococcus aureus in 41% of cases. The median duration between surgery and abscess was 1.5 months, underscoring the need for postoperative vigilance. To address the risk of postoperative cerebrospinal fluid leakage, proposed strategies include autologous bone plate placement. A modified approach in cases with disrupted barriers demonstrated success in 183 patients, cautioning against surgical injury, especially anticipating postoperative irradiation, reflecting ongoing efforts towards refined neurosurgical practices. This case underscores the atypical nature of his complications, diagnostic complexities, and multifaceted management of bacterial meningitis, including the rarity of cerebritis and the development of a CSF leak. This leak long after a neurosurgical procedure. It emphasizes the importance of interdisciplinary collaboration and prompt recognition in managing not only the common but also the atypical clinical presentations of this life-threatening condition. Further investigations are required to understand the long-term implications of intracranial complications in patients with bacterial meningitis.

Resident Poster # 006 Category: Clinical Vignette

Residency Program: Ascension Providence
Presenter: Jessica LaVoie
Additional Authors: Everett Woods, MD; Nabeel Shabo, MD

Idiopathic Communicating Hydrocephalus: Not all Ventricles are Created Equally

Hydrocephalus occurs when there is abnormal accumulation of cerebrospinal fluid in the ventricles, or spaces within the brain. This can be due to either overproduction, obstruction, or problems with reabsorption of CSF fluid. There are two classes of NPH, Idiopathic or primary NPH, where identifiable cause is found, and symptomatic

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 007 Category: Clinical Vignette

Residency Program: Ascension Providence
Presenter: Keyur Patel
Additional Authors: Alex Saunders, DO; Frank Shay

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster #: 009 Category: Clinical Vignette

Residency Program: Ascension Providence
Presenter: Frank Shaya
Additional Authors: Jessica LaVoie, DO1; Jyotsna Mary George, MD1; Keyur Patel MD1; David Putt, DO MBA1; Saif Affas, MD2

Mycoplasma Pneumoniae Presenting with Cavitary Lung Lesions: A Case Report

Introduction:

Mycoplasma pneumoniae is a common cause of atypical community-acquired pneumonia (CAP), responsible for up to 30% of cases [1]. It typically causes mild, self-limited illness. However, in rare cases it can escalate into severe manifestations like cavitary lesions [2]. We present a case of M. pneumoniae pneumonia with multiple cavitary lesions refractory to broad-spectrum antibacterial and antifungal treatment.

Case Presentation:

A 36-year-old woman with pertinent medical history of alcohol and tobacco use disorder presented for one week of worsening respiratory symptoms and fever. Exam showed tachycardia, tachypnea, and fever. Imaging revealed lung masses and cavitary lesions. She was admitted for atypical pneumonia and started on IV antibiotics. Despite escalating treatment, she deteriorated, requiring intensive care for hypoxemic respiratory failure. Serial imaging showed enlarging lung cavities. Bronchoalveolar lavage cultures were negative but pneumococcal antigen urine was positive. After an extensive viral, fungal, and bacterial infectious workup, ultimately Mycoplasma IgM was positive. With antibiotic adjustment including azithromycin, she clinically improved. Patient was ultimately discharged in stable condition and sent home on a 9-day PO steroid taper and oral antibiotic therapy.

Discussion:

Mycoplasma pneumoniae is a common bacterial cause of upper and lower respiratory tract infections, infecting 1% of the United States population annually. It can progress to community acquired pneumonia, which is typically atypical in nature with patients experiencing mild systemic symptoms. This report illustrates a rare presentation of severe M. pneumoniae CAP causing cavitary lesions refractory to broad-spectrum antibacterial and antifungal therapy.

Mycoplasma pneumoniae is a short rod with no cell wall, which makes the pathogen not visible on Gram Stain. It is not routinely cultured because of the special culture media it requires and the length of time required for growth (7-21 days) [3]. M. pneumoniae can also not be differentiated from other causes of atypical pneumonia such as Chlamydia and Leigenalla through clinical or radiological findings. If required, PCR can be used for rapid diagnosis [4]. Cold agglutinin test can also be carried out to support clinical diagnosis of M. pneumoniae.

Our patient was found to have a positive M. pneumoniae IgM, which is indicative of acute infection. Despite the positive pneumococcal urine antigen, no growth resulted from the bronchoalveolar lavage samples that were cultured, making streptococcus pneumoniae the unlikely causative organism. Mycoplasma PCR was also performed and the result was negative. However, results may have been affected by the patient being started on antibiotics promptly in the emergency department due to meeting sepsis criteria, i.e., antibiotics were initiated before samples for PCR were collected. Overall, this would indicate that mycoplasma pneumoniae was the most likely cause of the infectious process.

M. pneumoniae should be considered as a potential etiology of cavitary pneumonia after common bacterial and fungal etiologies have been reasonably excluded. Macrolide therapy led to clinical improvement in this patient, highlighting the importance of empiric therapy for atypical organisms in severe CAP even with negative cultures.

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 010 Category: Clinical Vignette

Residency Program: Ascension Providence
Presenter: Everett Woods
Additional Authors: Medhat Chowdhury (Cardiology Fellow), Dr. Souheil Saba (Attending)

Detection of Air Embolism on Transesophageal Echocardiography during Placement of a Left Atrial Appendage Occluder Device.

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 012 Category: Clinical Vignette

Residency Program: Ascension St. John
Presenter: Bhuwan Bhatta
Additional Authors: Archita Chandra ,Dr. Joel Topf, Dr. Hemapriya Gopalreddy,

Abstract for Bismuth Salicylate Toxicity Case

Abstract for Bismuth Salicylate Toxicity Case

Introduction: About 40% of the adult U.S. population uses over-the-counter antacids. Among them, bismuth subsalicylate is the most common, comprising 60% of the market. Each tablespoon (15 ml) of bismuth subsalicylate contains 130 mg of salicylate, or 302 mg per recommended dose. Original Pepto-Bismol is sold in sizes ranging from 118 ml to 473 ml. One large bottle of bismuth subsalicylate roughly contains 4 grams of salicylate. It requires only one and a half large bottles of bismuth-subsalicylate, 6.5 g, to cause acute salicylate toxicity. Salicylate disrupts the mitochondrial oxidative phosphorylation, inhibiting Tricarboxylic Acid Cycle and mitochondrial ATP production leading to lactic acidosis and accumulation of ketones causing increased anion gap metabolic acidosis.

Objective: demonstrate how bismuth subsalicylate overuse/misuse can precipitate salicylate toxicity.

Case: A 64-year-old male presented to the emergency department with a complaint of new-onset, progressively worsening nausea, bilateral tinnitus, and an episode of melanotic stools. He reported taking one and a half bottles of bismuth subsalicylate (estimated dose of 700 ml) to self-medicate for nausea, after which he developed tinnitus and the nausea worsened. He further reported taking two naproxen sodium tablets per day for 2 months to address post-exercise muscle pain. The physical exam was significant for decreased bilateral hearing and a black tongue. Initial labs showed salicylate levels of 34.0 mg/dL. Venous blood gas showed a pH of 7.39, PCO2 of 39mmHg, PO2 of 44mmHg, HCO3 of 23 mmol/L. He was started on a bicarbonate drip to alkalinize the urine. Symptoms improved. There was no indication for dialysis. Follow-up labs showed salicylate levels of 20.1 mg/dL; venous blood gas pH of 7.52, PCO2 of 32mmHg, PO2 of 65mmHg, HCO3 of 25mmol/L.

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 013 Category: Clinical Vignette

Residency Program: Ascension St. John
Presenter: Samer Dubaisi
Additional Authors: Sara Shaban MD, Ramtej Atluri DO, Philip Vendittelli DO

Beauty and the QT

Introduction:

Sudden Cardiac Death (SCD) is a complex, multifaceted phenomenon that can strike at any stage of life. On

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 015 Category: Clinical Vignette

Residency Program: Ascension St. John
Presenter: Yaqian Liao
Additional Authors: Heather Henderson <heather.henderson2@ascension.org>. Ahmad Sater <ahmad.sater@ascension.org>

A case of Streptococcus sanguinis endocarditis mimicking ANCA-associated vasculitis

Introduction:

The presentation of bacterial endocarditis may vary and mimic rheumatic disease. ANCA-positive bacterial endocarditis constitutes a special subgroup. ANCA positivity was detected in 4-31% of infectious endocarditis (IE), with 79% ANCA/PR3, 11% ANCA/MPO and 8% double positive. Here we present a male with a vasculitic rash later diagnosed with streptococcus sanguinis IE. He has positive ANCA and biopsy evidence of leukocytoclastic vasculitis (LCV).

Case:

A 50-year old male presented with rash. Medical history significant for ESRD 2/2 FSGS, chronic diastolic heart failure, aortic insufficiency and essential hypertension.

He initially developed a rash on left hand 2 days prior to presentation spreading to the upper left arm. He noticed continued spread to bilaterally lower extremities with mild itchiness. He denies recent vaccinations, new medications, or dental procedures. He started hemodialysis a month prior by left arm AV graft.

ROS: Intermittent sharp burning chest pain for a month alleviated by leaning forward (he finished 5 days of prednisone without improvement). Dark urine and fatigue. Otherwise negative.

Physical exam on presentation: blood pressure 131/55, HR 103, RR 22, afebrile, 100% saturation on room air. Petechiae and purpura with central scabbing over the left arm, bilateral feet, cheeks and nose tip. Pitting edema on left lower leg and left arm.

Pertinent labs: WBC 7.84 K/mcL, hemoglobin 8.2 g/dL, platelet 154 K/mcL. Urinalysis showed moderate blood with 20 RBC/HPF, low C4 (10 mg/dL), elevated CRP (38 mg/L).

Autoimmune vasculitis was initially suspected with plan to initiate steroids. Extensive rheumatic workup only positive for c-ANCA (1: 80). However, TTE completed on day 2 to evaluate pericardial effusion showed worse aortic regurgitation, small to medium pericardial effusion and a new multilobulated mass attached to the aortic valve later confirmed by TEE. Blood culture obtained grew Streptococcus sanguinis. Ceftriaxone was started on day 3.

On day 9, he underwent aortic valve replacement surgery. Repeat blood cultures were negative. Patient was discharged on day 17 to complete a 6-week course of ceftriaxone, with resolution of edema and near resolution of rash.

Discussion:

SBE may present with positive ANCA and LCV. In a study among 110 IE, prior biopsy-proven cutaneous LCV was found in 4. Reversely, retrospective study of 138 biopsy-proven cutaneous LCV, IE was identified as the culprit in 3.

A few clues in our patient may help guide us towards SBE. His risk factors include aortic valve insufficiency, recent hemodialysis initiation and recent fistulogram for AV graft malfunction. Part of his picture represents an overlap between rheumatic disease and IE including pericardial effusion, hematuria, fatigue, intermittent fever and LCV.

Conclusion:

Patients with SBE may present with LCV and test positive for ANCA. Empirical immunosuppression may improve ANCA vasculitis

2024 Michigan Chapter ACP and SHM Michigan Chapter

2024 Michigan Chapter ACP and SHM Michigan Chapter

2024

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 020 Category: Clinical Vignette

Residency Program: Ascension St. John
Presenter: Waleed Rehman
Additional Authors: Dr. Salem Sayar

Unleashing the Thunder: A Case of Electrical Storm

Introduction:

Electrical storm is a life-threatening syndrome that is defined by 3 or more sustained episodes of ventricular tachycardia, ventricular fibrillation, or appropriate shocks from an implantable cardioverter-defibrillator (ICD) within 24 hours. Electrical storm can manifest itself during acute myocardial infarction and in patients who have structural heart disease, or an inherited arrhythmic syndrome. Today we present a case of electrical storm in a patient with a known history of infiltrative cardiomyopathy due to sarcoidosis.

Case Report:

A 60-year-old woman with a medical history significant for atrial fibrillation on Eliquis, infiltrative cardiomyopathy due to sarcoidosis with an ICD in place, hyperlipidemia, and hypertension presented to the emergency department after receiving a shock from her defibrillator.

The patient reported a sudden onset of uneasiness in her upper epigastric region, escalating until the defibrillator shock prompted her visit to the emergency department. On arrival, she experienced six rounds of sustained monomorphic ventricular tachycardia, remaining hemodynamically stable but expressing discomfort characterized by heaviness and esophageal contractions. Intravenous administration of amiodarone, lidocaine, and metoprolol tartrate initially alleviated symptoms. She was subsequently transferred to the cardiovascular intensive care unit (CVICU).

In the CVICU, the patient continued to experience episodes of sustained ventricular tachycardia. She was started on a combination of a lidocaine drip, esmolol drip, and continued on amiodarone. This regimen successfully restored normal sinus rhythm. Upon discharge, she was prescribed oral amiodarone (400 mg twice daily), toprol-XL (75 mg twice daily), and mexiletine (150 mg three times daily).

Discussion:

The origin of electrical storms spans a spectrum, from myocardial ischemia to infiltrative cardiomyopathy, as observed in our patient. Decoding the morphology of the QRS complex serves as a diagnostic clue, with monomorphic ventricular tachycardias typically linked to structural heart diseases, while polymorphic ventricular tachycardias are associated with myocardial ischemia. Symptoms may manifest as palpitations, dizziness, and

Resident Poster # 021 Category: Clinical Vignette

Residency Program: Ascension St. John
Presenter: Ahmad Sater
Additional Authors: Hussein Gharib MD; Maxwell Wheaton, BMS; Raghavendra Kamath, MD

A Rare Case of Computer Duster Inhalant Abuse Complicated by Thoracic Syringomyelia

Introduction: Computer duster is a widely available aerosolized can used for cleaning and contains Nitrous Oxide, Difluoroethane, Alkyl Nitrates and various other chemicals. Nitrous Oxide and amyl nitrates have become commonly abused insufflated drugs recently. Nitrous oxide irreversibly oxidizes the cobalt ion of cobalamin (Vitamin B12) depleting the body's vitamin B12 altering myelin formation. Nitrous oxide use is not common, but

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 022 Category: Clinical Vignette

Residency Program: Ascension St. John
Presenter: Beth Schwartz
Additional Authors: Ismail Malik MD, Paul Kudla MD, Leonard Johnson, MD

A Case of Sparganosis in Breast Tissue

Introduction:

Sparganosis is a parasitic infection, caused by Spirometra tapeworm larvae, which usually presents in humans as a subcutaneous nodule . Infection in the United States is rare, with the majority of cases reported in east and southeast Asia. As of 1996, approximately 62 cases of sparganosis were reported in the United States. We highlight the importance of awareness of parasitic infections in the differential diagnosis of an enlarging breast mass.

Case:

A 69-year-old male with hypothyroidism presented for a right breast lump. He noticed it three years ago [1c12 7(9)] and since then it has varied in size, growing and shrinking unpredictably. He stated it had become increasingly pruritic over the last 12 months and endorsed occasional tenderness. He denied nipple discharge. He denied ingestion of uncooked seafood, raw frogs, tadpoles, snakes, or residence in Southeast Asia. However, he endorsed that as a child he played with tadpoles and frogs. He lived in rural Asia and drank well water until adolescence. Vitals were unremarkable and exam showed a firm, tender right chest wall lump without induration or drainage. Complete blood count showed a normal white blood cell count without eosinophilia. Ultrasound and mammogram

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster #: 024 Category: Clinical Vignette

Residency Program: Ascension St. John
Presenter: Abdullah Shaik
Additional Authors: Abdullah Shaik, MD, Waleed Rehman, MD, Adam Zakkar, DO, Ahmed Elshaer, MD, Michael Kern, MD

A Novel Case of Reverse Takotsubo Cardiomyopathy Following Gastroenteritis

Takotsubo syndrome (TTS) also known as stress cardiomyopathy is a condition that causes a temporary change in the heart's contractility. Generally, involves apical akinesis, basal hypercontractility, and regional wall motion abnormalities beyond single vascular distribution. Resembles acute coronary syndrome (ACS) and is hypothesized as triggered by emotional or physical stress, often affecting postmenopausal women. Reverse Takotsubo Syndrome - rTTS is a rare variant of TTS and is characterized by apical akinesis and basal segment depression. We present the case of a 72-year-old male with a history of prostate cancer, diabetes mellitus, hypertension, and hyperlipidemia who developed rTTS following symptoms of gastroenteritis. Vitals upon arrival showed a heart rate of 81 bpm, blood pressure 164/94 mmHg, and respiratory rate of 16. The patient presented with fatigue, nausea, vomiting, subjective fever/chills, and abdominal pain. Initial assessments in the emergency department revealed leukocytosis (22.92), elevated D-dimer (5270), and elevated Troponin levels (0.11) with a repeat troponin of (0.11). EKG showed sinus rhythm, with no ST segment elevation or depression. Furthermore, CK was elevated (309), along with a lactate of as (1.8). Chest x-ray and CTA chest were unremarkable ruling out pulmonary

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 026 Category: Clinical Vignette

Residency Program: Beaumont Health Dearborn
Presenter: Khurram Arshad
Additional Authors: Zaid Mahameed, Khurram Arshad, Rabia Latif, Ruaa Elteriefi

Yazan Alamro

From Collapse to Recovery: Thiamine Intervention in Cardiac Beriberi.

Introduction:

Thiamine deficiency, historically known as Beriberi, poses diagnostic dilemmas due to its potential occurrence without overt malnutrition or alcohol dependence. This case report explores the intricate dynamics of cardio-circulatory collapse coupled with pericardial effusion, unveiling an underlying thiamine deficiency.

Case Description:

A 68-year-old male with a medical history of hypothyroidism, heart failure, and chronic mild pericardial effusion presented with worsening dyspnea, leg edema, and hypotension. The patient's physical evaluation disclosed notable findings, including marked obesity, distant heart sounds, an audible S4 sound, and swelling in the lower extremities. Laboratory results revealed chronic anemia. Brain natriuretic peptide (BNP) was mildly elevated at 286 pg/mL, accompanied by severe hypoalbuminemia (1.6 g/dL) and prealbumin levels (6 mg/Dl). The coagulation panel and cardiac ischemia workup were within normal limits.

The patient underwent surgical pericardial effusion drainage with a pericardiocentesis drain. Despite the intervention, the patient developed hypotension unresponsive to fluid resuscitation, prompting a septic workup and initiation of vasopressors. We init

2024

2024 Michigan Chapt BT /F1 t mAC202

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 029 Category: Clinical Vignette

Residency Program: Beaumont Health Dearborn
Presenter: Tabssum Chauhan
Additional Authors:

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 030 Category: Clinical Vignette

Residency Program: Beaumont Health Dearborn
Presenter: Antoine Egbe
Additional Authors: Mariam Jamil, Hussein Gaith, Wafa Ahmed, Ali Mozaffari, John Rajiv, Essa Kadiri

Navigating Recurrent Acute In-Stent-thrombosis: Insights from a Single Admission Case Study

In-stent thrombosis (ISR) is one of the dreaded complications of percutaneous coronary angioplasty. Despite marked reductions in ISR since the transition from bare metal stents to drug-eluting stents, ISR still occurs at a rate of 1%-2 % per year. We present the case of a 64-year-old Caucasian male who was admitted for staged angioplasty after a left heart catheterization had shown he had significant stenosis to the LAD and mid ramus intermedius. He had DES placed to the mid-LAD and to the mid-ramus intermedius. Despite being placed on dual antiplatelet therapy after the stent placement, 6 hours later he complained of chest pain and EKG showed new ST elevations in leads 1, AVl, and V2-V3. Repeat LHC showed thrombosis to recently placed LAD and mid-ramus intermedius stents so a new set of stents was placed and the patient was loaded with eptifibatide and then restarted on dual antiplatelet therapy after procedure.

However, only 48 hours later the patient started complaining of chest pain again. EKG showed new ischemic changes and a high sensitivity troponin of 5000. He was taken to the catheterization and LHC once again late stenosis of the previously stented LAD and mid-ramus intermedius. He underwent percutaneous coronary angioplasty to these stents. Post-LHC he was started on prasugrel and aspirin instead of the 2 other times when he was on clopidogrel and aspirin then ticagrelor and aspirin.

In addition to the dual antiplatelet therapy, he was also started on a Factor X inhibitor, rivaroxaban. Our hypercoagulability investigations did not yield any positive result, however we found out the patient was on testosterone replacement.

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 032 Category: Clinical Vignette

Residency Program: Beaumont Hea Category:

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 033 Category: Clinical Vignette

Residency Program: Central Michigan University
Presenter: Denise Mourad
Additional Authors: Mourad, Denise. Petreska, Natasa. Letterio, Kenneth. Crowlton, Nicholas. Mohammed, Adil. Jarad, John. Mathew, George. Haddad, Nicholas.

Wased Al-Khateeb

Unveiling a New Flow Enigma: West Nile Virus and the Intriguing Threat of Hypercoagulability

West Nile Virus (WNV) is a single-stranded RNA virus, a part of the Flaviviridae family. It is transmitted to humans by mosquito bites. Clinical presentations range from asymptomatic to severe infections such as encephalitis. A rarely reported outcome of WNV infection is hypercoagulability. We present the case of a 72-year-old male patient who presented with a complaint of altered mental status (AMS), secondary to WNV infection, and an incidentally found superior sagittal sinus thrombosis.

Our patient has a history of hypertension, type 1 diabetes, and bipolar disorder. He was found by his family to be minimally responsive and was brought to the emergency department. On presentation, his vital signs were within normal limits, and due to his AMS, he was promptly initiated on antibiotics for suspected meningitis, together with anticonvulsants for possible status epilepticus. He was admitted to the ICU for encephalopathy. Workup included a complete blood count (CBC) showing mild leukocytosis, and a complete metabolic panel (CMP), which was noncontributory. Lactic acid level was within normal limits, urine drug screen was positive only for Cannabinoids. Electroencephalogram (EEG) ruled out seizures. Initial brain imaging included a computed tomography (CT), and subsequently a magnetic resonance imaging (MRI) both demonstrating age-related changes and small prior ischemic events. His antibiotics were de-escalated, and his antiseizure therapy was discontinued.

During his hospitalization, this patient demonstrated some clinical improvement with spontaneous eye opening, but his mentation remained altered. Lumbar puncture (LP) and repeat brain imaging were ordered. On day seven of hospitalization, a brain magnetic resonance angiogram (MRA) uncovered a filling defect of the anterior portion of the superior sagittal sinus. This was confirmed with a repeat MRI, and computed tomography angiogram (CTA) to be an anterior superior sagittal sinus thrombosis. The cerebral spinal fluid (CSF) analysis showed 11 WBC

Resident Poster # 034 Category: Clinical Vignette

Residency Program: Central Michigan University
Presenter: Keith Bouffard
Additional Authors: Paritharsh Ghantasala MD, John Jarad MD, Sayed Osama MD

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 035 Category: Clinical Vignette

Residency Program: Central Michigan University
Presenter: Kinga Budnicka
Additional Authors: Mathew Kunz, MD., Jibran Ajaz, MD., Robert Kasemodel, MD., Jonathan Rene, MD.

A Rare Case of Hypomyopathic Dermatomyositis

Dermatomyositis (DM) is both a complex and rare inflammatory myopathy affecting 1 in every 100,000 individuals, most commonly in women ages 40-60. Classically, DM presents with progressive proximal muscle weakness in conjunction with characteristic dermatological findings, which are highly disabling. However, it is important to note that the timeline of these symptoms may present differently in a subset of patients. I will present a very rare case of hypomyopathic dermatomyositis in a young 26 year old female.

This is the case of a 25 year old female with unremarkable past medical history that presented to the rheumatology clinic for a positive ANA result. She mentioned that the ANA was ordered after a thyroid nodule was found by her PCP, and since then, she started having some noticeable fatigue. Additionally, she has had both weight and hair loss. On further prompting, the patient noted a skin rash on her hands, nodular in nature on a few of her metacarpal and proximal interphalangeal joints. She confirmed that she has had a red rash across her cheeks before and that her PCP believes she likely has Lupus. She has no family history of cancers however she does have an uncle with Lupus. The physical exam was positive for mild Gottron's hands, erythematous scaly papules over 2 knuckles, periungual erythema and an erythematous upper back. The patient was asked to complete some rheumatological labs, which came back exhibiting an ANA of 1:1280, elevated TPO and iron deficient Thalassemia. TSH, ESR were normal. DsDNA, ENA, anti-Smith, Anti-Jo1, alpha 1 antitrypsin, ssDNA, SSA, SSB were all negative. There were multiple differentials at the time ranging from autoimmune thyroiditis, dermatomyositis, anti-synthetase syndrome, SLE and others. After 6 months, the patient re-presented to the clinic due to worsening symptoms, including muscle weakness such as difficulty gripping objects and some intermittent numbness in the hands. EMG demonstrated electrodiagnostic evidence of myositis present proximally and distally. Right thigh biopsy was performed and sent for analysis showing myositis. Additional labs completed showed CK of 5719, Aldolase of 36.6, AST 239, ALT 129. Anti-centromere, Anti-chromatin, Anti-Ribosomal P, Anti SCL 70, and Anti RNP Abodies came back negative. Patient was started on a high dose steroid taper, and placed on immunosuppressive medications including Methotrexate and Plaquenil. Due to the association of DM with cancer, appropriate cancer screening was performed as well. Once her disease was more stabilized, the patient was referred for low grade physical therapy. She was officially diagnosed with Hypomyopathic DM, due to the late presentation of muscle symptoms, stirring away from the classical presentation of the disease.

This case demonstrates the importance of a concrete history and physical. The scattered timeline of cutaneous and muscle symptoms and the plethora of symptoms in the following case is infrequently encountered, making it a formidable diagnostic challenge. Recognizing this uncommon inflammatory myopathy is critical in initiating early appropriate therapy, due to its debilitating symptoms and cardiovascular, pulmonary and/or oncologic complications if left untreated.

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters
Resident Poster # 037 Category: Clinical Vignette

Residency Program: Central Michigan University
Presenter: Natasa Petreska
Additional Authors: Dr. Nicholas Haddad, Dr. Kelly Blouffard, Dr. Wasef Al Khateeb, Dr. Matthew Kunz

Dr. John Jarad (PGY2), Dr. Natasa Petreska (PGY1), Dr. Nicholas Haddad, FACP, FIDSA, CNE, AAHIVS

The New ART: Parenteral Antiretroviral Therapy Achieving Viral Suppression

Introduction:

Daily oral antiretroviral therapy (ART) is the standard of care for treatment of HIV. It is composed of three drugs, taken once daily, which reliably achieves durable viral suppression when taken with strict compliance. Cabotegravir and rilpivirine have been approved as a two-drug long-acting injectable antiretroviral therapy (LAIART). The current indicated use of LAIART is for maintenance therapy in patients who have already achieved virologic suppression (HIV-1 RNA <50 copies/mL) on stable ART for at least 3 months without previous treatment failures. LAIART is not presently approved as a first-line induction ART. This new LAIART represents a convenient treatment modality in patients who prefer once monthly or bimonthly (every 2 months) injections in clinic as opposed to daily oral therapy. There are currently no large studies demonstrating efficacy in patients with high viral loads. Recent reports are emerging suggesting they may also induce durable viral suppression. Control of HIV viral load not only improves patient mortality and quality of life, but also carries public health benefits for the prevention of the spread of HIV.

Case Presentation:

A 24-year-old male who achieved prompt viral suppression with the unindicated use of LAIART. The patient had congenitally acquired HIV infection, and neurocognitive deficit. While under the care of his guardian (grandmother), he was virologically controlled with Elvitegravir / Cobicistat / Emtricitabine / Tenofovir, the daily intake of which she was supervising (CD4 at 561 cells/mL, HIV undetected). When the patient turned 18, he left home, and was lost to follow up with our infectious disease clinic. His family later reassumed guardianship at 23 years of age. He re-established care in the infectious disease clinic at their behest and was found to have developed AIDS. His CD4 cell count of 89 cell/uL, and an HIV1 viral load at 2,670,000 copies/microL. His previous enteral therapy was re-initiated. The patient initially reported compliance (VL down to 25,139 cells/microL) but within 4 months, his CD4 cell count had decreased to 35 cells/uL and his HIV1 viral load had increased to 394,946 copies/mL. Multiple attempts at garnering compliance failed. It was decided through joint decision making that initiating LAIART with Cabotegravir and rilpirivine as monthly IM therapy would increase the patient's adherence to therapy and may improve viral control. The patient was himself agreeable and over the next 6 months presented to the clinic for monthly injections. Subsequently, his CD4 cell count improved to 93 cells/uL, and viral load was detectable but less than 20 copies/mL. 6 months after achieving suppression, the patient continued to have an undetectable viral load.

Discussion:

HIV infection is often co-morbid with many social barriers that prevent the effectiveness of oral therapies to induce remission. LAIART overcomes many of these issues, but there are currently no recommendations for this indication. Case reports suggest that LAIART may be a useful alternative therapy and further studies to explore the effectiveness of LAIART for inducing durable viral suppression should be pursued.

Resident Poster # 038 Category: Clinical Vignette

Residency Program: Central Michigan University

2024 Michigan Chapter ACP and SHM Michigan Chapter

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 040 Category: Clinical Vignette

Residency Program: Central Michigan University
Presenter: Denise Mourad
Additional Authors: Jaspreet Nannar, John Jarad, Dr. Vivek Variar

Pheochromocytoma recurrence; is lifelong surveillance necessary?

Introduction

Pheochromocytomas are rare neuroendocrine tumors originating from the adrenal glands. They are typically identified during work-up of resistant hypertension usually associated with episodes of hyperhidrosis and headache. Untreated, pheochromocytoma can result in hypertensive crises leading to strokes, malignant hypertension, myocardial ischemia, and heart failure which could often be lethal. Surgical resection is mostly curative. Surveillance after surgery is needed to detect recurrence. Studies have shown that the mean time to recurrence following a resected sporadic pheochromocytoma is about 49.4 months with a recurrence rate of approximately 3% (1). It is possible that recurrence may present with metastatic disease. However, there are no specific recommendations regarding the duration for surveillance after resection.

Case Presentation

A 47-year-old African American female underwent right sided adrenalectomy for pheochromocytoma. During the first two years of postoperative follow up, she was asymptomatic, and her catecholamine levels were normal. Four years after initial surgery, her blood pressure was elevated during an office visit with primary care physician. Workup revealed elevated plasma metanephrines level by >30% and she was referred to our clinic. On evaluation, there was concern for pheochromocytoma recurrence. A computed tomography (CT) scan of the abdomen showed a mass extending into the porta hepatis region, suggestive of recurrent pheochromocytoma. A PET CT per adrenal protocol confirmed a well-defined somatostatin receptor 2 positive soft tissue mass in the right retroperitoneum surrounding the superior pole of the right kidney, which extended anteriorly to the porta hepatis, and postero-medially to the perihepatic space in addition to multiple nodular soft tissue densities with increased radiotracer uptake in the right paracolic gutter suggestive of metastatic deposits. To decide the next best step for care, she was referred to a tertiary care center for opinion. After a tumor board review, she was offered four options which included experimental therapy with Belzutifan vs chemotherapy with capecitabine and temozolomide vs lutetium dotatate vs no intervention. She eventually was started on peptide receptor radionuclide therapy (PRRT) using lutetium dotatate.

Conclusion

Pheochromocytoma can be a fatal disease if not identified and treated promptly. Patients can become completely

Resident Poster #

2024 Michigan Chapter ACP and SHM Michigan Chapter

2024 Michigan Chapter ACP and SHM Michigan Chapter

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 046 Category: Clinical Vignette

Residency Program: Corewell Health William Beaumont University Hospital
Presenter: Nitya Batra
Additional Authors: Nitya Batra 1 , Atulya Aman Khosla 1 , Mohammad Mudassir Chishti 1 , Vikash Jaiswal 2 , Ashbita Pokharel 1

A rare case of primary sclerosing pneumocytoma in a healthy female

Introduction: Pulmonary sclerosing pneumocytoma is a rare group of benign pulmonary neoplasms that usually affects older adults over 50 years of age. This neoplasm is more commonly seen in women, with female to male ratio of 5:1 and is predominantly seen in Asian ethnicities. These tumors originate from type II alveolar pneumocytes and are hence classified as adenomaunder the 2021 World Health Organization (WHO) classification of tumors . They account for 5-3 percent of total benign lung tumors. One of the striking aspects of PSP is its clinical presentation, or rather the lack thereof in many cases. It is often discovered incidentally during routine chest imaging, such as chest X

Resident Poster # 047 Category: Clinical Vignette

Resident Poster # 048

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 054 Category: Clinical Vignette

Residency Program: Detroit / Wayne County Authority Health
Presenter: Ayman Mahdi
Additional Authors: Nwonukwuru Amadi, MD (Resident), Rishab Rajendra Prabhu, MD (Resident), Abdulkader Hmidan Simsam, MD (Resident), Cecilia Cosma (Faculty), Dr. Ruhi Mahayni (Faculty)

Adenovirus and Herpes Simplex Virus 1 co-infection leading to acute respiratory distress syndrome (ARDS) in a latent mycobacterium tuberculosis, immunocompetent, patient

Introduction:

Adenovirus and Herpes Simplex Virus 1 (HSV-1) typically cause self-limiting upper and lower respiratory tract infections. However, in immunocompromised individuals, severe complications such as pneumonia can arise, leading to acute respiratory distress syndrome (ARDS). Management typically requires prompt intubation and mechanical ventilation for survival. This case report details a rare occurrence of co-acquired adenovirus and HSV-1 pneumonia in an immunocompetent adult, resulting in ARDS.

Clinical Vignette:

A 40-year-old Hispanic female, without a known medical history, presents to the hospital with sudden left chest pain, shortness of breath, and hemoptysis lasting one day. The pain, rated 10/10, began on the left chest side and radiated to the center back, exacerbated by deep breaths and upper body movement. Over-the-counter ibuprofen provided no relief, prompting hospital evaluation. In the ED, vital signs revealed a temperature of 98.2°F, heart rate of 116, respiratory rate of 27, blood pressure of 135/91, and SpO2 of 95% on 6L nasal cannula.

Initial labs were remarkable for sodium 133, potassium 3.4, chloride 99, BUN 14, creatinine 0.66, lactate 1.4, troponin 6, WBC 13.7, hemoglobin 12.7, hematocrit 38.4, and platelet 280. Urinalysis was unremarkable. COVID test was negative. CT Angiogram Chest revealed severe multifocal pneumonia but no pulmonary embolus. Before transferring to medicine services, the patient received ceftriaxone, azithromycin, and vancomycin. Overnight, the patient's condition worsened, necessitating Vapotherm oxygen therapy and eventually BiPAP. Imaging and blood gases confirmed moderate ARDS, with ABG showing pH 7.46, pCO2 36.4, PO2 49, and HCO3 25.7.

ICU was consulted for continuous BiPAP use. She continued to worsen and was intubated overnight. She was

r 1 0612 G 36 G [(a)-20(nd S)10(pO)-12(2)7(o)3(f)5(9)8(5)7(%)-9(o)3(n 6)8(L)-4(na otcb)11(2)-16(8)74, fngeSu 1 0

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 055 Category: Clinical Vignette

Residency Program: Detroit Medical Center - Huron Valley - Sinai Hospital
Presenter: Conrad Chrabol
Additional Authors: Karina Torres, Nzube Ekpunobi, Narod Kalaijean

A post-transplantation puzzle: a case of recurrent fevers and lymphadenopathy

Lymphadenopathy, often first noted on a physical exam, can create a diagnostic dilemma when an obvious cause is not apparent from the medical history. Imaging studies such as ultrasound are frequently performed for further lymph node characterization, or to distinguish between local and generalized involvement. Diagnosis can ultimately require a biopsy to reveal specific pathologic features or staining patterns. Lymphadenopathy often coexists with non-specific symptoms such as fevers, myalgias, and fatigue. The differential diagnosis can usually be grouped into infectious, cancerous, lymphoproliferative, and immunologic causes. In the setting of immunosuppression, as in post-transplant recipients, a high index of suspicion must be kept for rare or unusual disease processes, especially if the initial work-up is non-revealing.

A 32-year-old male with a history of ESRD secondary to FSGS, s/p kidney transplant 10 years prior (maintained on tacrolimus and mycophenolate), with recent transplant rejection requiring reinitiation of dialysis, presented to the hospital due to worsening fatigue, fevers, shortness of breath, and episode of syncope. He had recently been discharged after an extended hospitalization for similar presenting symptoms. Routine labwork had shown a significantly elevated white blood cell count (neutrophil-dominant), anemia (thought to be mainly from renal disease), and thrombocytopenia, but were otherwise unrevealing. Patient's vitals were significant for labile-hypotensive blood pressure and recurrent fevers to the point of rigor, poorly responsive to antipyretics. Patient's exam was notable for a generalized pruritic maculopapular rash that was not responsive to topical steroids. Chest imaging revealed diffuse patchy airway disease thought to initially represent multifocal pneumonia, though with multiple enlarged thoracic lymph nodes. Abdominal imaging revealed hepatosplenomegaly. He was empirically started on broad spectrum IV antibiotics, without improvement. An extended infectious diagnostic panel was performed, which returned negative for bacterial, viral, and fungal pathogens. Bronchoalveolar lavage was likewise unrevealing. A bone marrow biopsy was negative for malignancy. In the absence of an identifiable infectious cause, and history of prolonged immunosuppression, the presumptive diagnosis was thought to be post-transplant lymphoproliferative disease. A lymph node biopsy was performed to confirm this diagnosis, which instead revealed a plasmablastic pattern of histologic features consistent with HHV-8 associated multicentric Castleman disease. Staining for latency-associated nuclear antigen (LANA) returned positive, confirming HHV-8 infection. Skin biopsy of the lesions was consistent with a neutrophilic dermatosis, not suggestive of Kaposi sarcoma (an angioproliferative disorder also mediated by HHV-8).

Castleman disease represents a rare disease of lymph nodes on the spectrum of lymphoproliferative disease and

Resident Posters

Resident Poster

Resident Poster # 059 Category: Clinical Vignette

Resident Poster # 060 Category:

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 062 Category: Clinical Vignette

Residency Program: Detroit Medical Center/Wayne State University
Presenter: Mohamed Alrayyashi
Additional Authors: Abdul Rasheed Bahar, Adi Zaclli, Pamila Sharma, Avneet Kaur Arora, Kamalakar Nerusu

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 063 Category: Clinical Vignette

Residency Program: Detroit Medical Center/Wayne State University
Presenter: Avneet Kaur Manjeet Singh Arora
Additional Authors: Mohamed Alrayyashi, Adi Zaclli, Kamren Huizenga, Vishakh Keri, Abdul Rasheed Bahar, Pamila Sharma, Lea M Monday

Disseminated Mycobacterium Abscessus with skin nodules and and septic arthritis following systemic immunosuppression in a patient with suspected primary immunodeficiency misdiagnosed with atypical sarcoidosis

Introduction:

Nontuberulous mycobacteria (NTM) are environmental pathogens found in water and soil which seldom cause disease in normal hosts but can present with a variety of clinical syndromes in immunocompromised patients. Mycobacterium abscesses is a rapidly growing NTM associated with a variety of infections ranging from pulmonary disease to cutaneous infection and dissemination in susceptile hosts. Here we present a patient who had been profoundly immunosuppressed by treatment for sarcoidosis not fitting typical diagnostic criteria and subsequnelty developed M.abscesses septicemia with dissemination to skin and joints.

Case Discussion:

A 21-year-old gentleman presented with fevers, painful skin nodules, and 35 pound weight loss of 12 months duration. He had been diagnosed with sarcoidosis based on retroperitoneal lymph node biopsy showing lymphoplasmacytic necrotizing inflammation suggestive of atypical granuloma which was never sent for culture. He worsened on several courses of steroids then 2 doses of infliximab. His father had died at age 26 due to pneumonia and had also been diagnosed with sarcoidosis but had never received a biopsy. Exam was significant for fever, tachycardia, micrognathia, atypical facies, and cachexia. There was diffuse palpable lymphadenlopathy, multiple purupic nodular lesions of the hands/fingers, hepatospelenomegaly, and a right knee effusion. Labs showed pancytopenia and blood cultures had M. abscesses in both standard and lysis-centrifugation bottles. Bone marrow biopsy was hypocellular (~15%) with depressed hematopoesis without granulomas or lymphohistiocytosis. HIV screening and an extensive rheumatology work up was nonrevealing. Knee arthrocentesis had 6200 nucleated cells (80% lymphocytes) with acid-fast bacilli on staining. He declined skin biopsy. Due to reticuloendothelial involvement, abnormal facies and family history of early death due to infection, inherited diseases strongly suspected that disseminated NTM infection due to an an inherited immunodeficiency and infliximab. Further immunosuppression was held and he

Tf 1 0 0 1 284.h70.8 s 0 g 0 (1)7(5)1sesattzive o fthy,

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 064 Category: Clinical Vignette

Residency Program: Detroit Medical Center/Wayne State University
Presenter: Abdul Rasheed Bahar
Additional Authors: Tushar Mishra, Mahmoud Othman

Aortic Valve Vegetation Due to Nonbacterial Thrombotic Endocarditis in a Patient with Antiphospholipid Antibody Syndrome

Introduction

Nonbacterial thrombotic endocarditis (NBTE) is a rare condition characterized by the development of sterile
Nc een-BDC q n Q q 0.02 6e De(t)10(2(E)-1792 re612 n-BD(t)10(moster)50(e Ucommonl-BD0.00(he d(t)10(t)10oc)-3(B)

Resident Poster # 066 Category: Clinical Vignette

Residency Program: Detroit Medical Center/Wayne State University
Presenter: Mariam (Gabby) Krikorian
Additional Authors: Michelle Malik, Ayman Salem, Abubaker Farah, Aws Polina

Resident Poster # 068 Category: Clinical Vignette

Residency Program: Detroit Medical Center/Wayne State University
Presenter: silvianovakova
Additional Authors: Muhammad Mushtaq DO, Zahia Esber, MD

Resident Poster #: 070 Category: Clinical Vignette

Residency Program: Detroit Medical Center/Wayne State University
Presenter: Sahalia Rashid
Additional Authors: Sahalia Rashid, MD, PhD., Tanushri Bushan, MD., Adam Qazi., Jessica Liang, MD., Brianna Schwarcz, MD., Rahul Vyas. MD.

Heparin-Triggered Inferior Epigastric Artery Bleeding Leading to a Remarkable Rectus Sheath Hematoma

Rectus sheath hematoma represents a rare clinical entity, exhibiting an incidence rate of 0.1-2%. The etiology of this hematoma remains largely elusive, contributing to its frequent misdiagnosis or underdiagnosis among physicians due to its infrequency. Trauma to the superior epigastric artery is promptly tamponaded by the rectus sheath. Conversely, injury to the inferior epigastric artery poses challenges in control, attributed to the absence of a posterior sheath.

The presented case involved a 72-year-old female with a myriad of comorbidities, including arthritis, hypertension, hyperlipidemia, diabetes mellitus, asthma, and sickle cell trait. Following admission for encephalopathy with concurrent COVID-19 infection, the patient's management included dexamethasone, supplemental oxygen, and subcutaneous heparin for DVT prophylaxis. Notably, she developed abdominal pain during her hospital stay vomiting CT Abdo/pelvis revealed a 6.8 cm rectus abdominis hematoma, suspected to be related to subcutaneous heparin injections. Despite the absence of acute surgical indications as noted by the general surgery team, the patient's clinical course took a critical turn with worsening encephalopathy, leukocytosis, elevated lactate, and hematologic abnormalities with a drop in hemoglobin from 8.1. A septic workup was initiated, and empirical broad-spectrum antibiotics were administered. Subsequently the patient deteriorated into shock, necessitating transfer to the MICU for further management. CTA Abdo/pelvis revealed an interval increase in the rectus sheath hematoma size to 12.4 cm, accompanied by contrast blushing indicative of active bleeding from the inferior epigastric artery. Additionally, hematoma and diffuse subcutaneous soft tissue suggested an extension from the rectus sheath hematoma. Interventional radiology was promptly consulted for emergent embolization, resulting in hemodynamic stabilization and subsequent improvement in hemoglobin levels.

The case underscores the importance of recognizing and promptly managing rectus sheath hematoma, particularly in patients with complex medical histories. The atypical course of this patient with potential sepsis and active bleeding, leading to shock underscores the multifaceted nature of this condition necessitating a multidisciplinary approach to optimal patient care.

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 072 Category: Clinical Vignette

Residency Program: Detroit Medical Center/Wayne State University
Presenter: Almustafa Kadem
Additional Authors: Mohamed Alrayyashi, Almustafa Kadem, Abdul Rasheed Bahar, Avneet Kaur Arora, Sajith Matthews

Pamila Sharma

A Case of Ibrutinib Withdrawal in a Patient with CLL

Introduction: Over the last decade, the introduction of targeted molecular therapy has become widely used in the treatment of specific cancers. Ibrutinib (brand name Imbruvica) is one such medication that works by inhibiting B cell proliferation by binding the Bruton's tyrosine kinase protein and preventing downstream signaling of this pathway. Although rare, withdrawal side effects have been reported in a small subset of patients using targeted molecular therapies and here we present a case of a patient with chronic lymphocytic leukemia who developed abnormal symptoms post medication cessation.

Case Description: Patient is a 59-year-old female with known medical history of chronic lymphocytic leukemia, diagnosed in 2020 following cervical lymph node biopsy, for which she has been treated with Ibrutinib 420mg daily. She presented to our emergency department with the acute onset of a severe headache, with associated fatigue, fever, and chills. To note, three days prior to presentation, patient had abruptly discontinued treatment with Ibrutinib at the recommendation of her hematologist as she had been experiencing m.178(ol)n-2(0 1 72.025 454.15 Tm 0

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 073 Category: Clinical Vignette

Residency Program: Detroit Medical Center/Wayne State University
Presenter: Nagasai Yalavarthi
Additional Authors: Yasir Alsaraf, M.D., Mohammed Najeeb Hallak, M.D.

A Rare Duo: An Atypical Case of Esophageal Adenocarcinoma in a patient with Lynch & Hereditary Breast and Ovarian Syndrome

Introduction: DNA mismatch repair enzymes are essential to maintain the integrity of DNA and prevent formation of cancer. Mutations in certain genes (MLH1, MSH2, MSH6, PMS2, EPCAM), and in (BRCA 1 & BRCA 2) are associated with Lynch and Hereditary Breast and Ovarian Syndrome (HBOC) respectively. These enzyme deficiencies increase the risk of gastrointestinal and genitourinary malignancies such as the traditionally associated

Resident Poster # 074 Category: Clinical Vignette

Residency Program: Detroit Medical Center/Wayne State University
Presenter: Mariam Zunnu Rain
Additional Authors: Mishita Goel, MD.,

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 075 Category: Clinical Vignette

Residency Program: Garden City Hospital
Presenter: Prathyusha Anumolu
Additional Authors: Aaron Gandhi DO, Ali Jishi MD, Sujata Kambhatla MD, Furqan Siddiqi MD

A Rare Case of Candida Pneumonia in the Intensive Care Unit

Introduction:

Pneumonia is a common respiratory tract disease and is one of the leading causes of mortality. Candida species rarely cause pneumonia, and candida pneumonia can be challenging to diagnose. C. glabrata is typically a nonpathogenic candida species and is present within normal respiratory flora. In this case report, C. glabrata pneumonia was diagnosed in a patient who is immunocompromised.

Case description:

A 64-year-old female with a history of lupus, scleroderma, thyroid cancer s/p resection, dilated esophagus, nicotine dependence, hyperlipidemia, hypothyroidism, and GERD initially presented to the ED due to shortness of breath and hemoptysis and was treated with azithromycin without improvement. Her CT findings were consistent with mucus plugging consistent with aspiration and post obstructive pneumonia on the left. Multifocal consolidations were observed bilaterally along with distended esophagus. Due to increased oxygen requirements, she was intubated and transferred to the ICU. Flexible bronchoscopy was performed which revealed mucus and

2024 Michigan Chapter ACP and SHM Michigan Chapter

2024

Resident Poster # 080 Category: Clinical Vignette

Residency Program: Garden City Hospital
Presenter: Mehwish Zeb
Additional Authors: Karishma Doshi, Amin Pasha, Sujata Kambhatla

Right thalamic abscess in a Caucasian man with rheumatoid arthritis, an intriguing case report.

Introduction:-

A brain abscess is a localized collection of pus within the brain tissue. Approximately 1500-2000 cases are presented annually in the United States. A thalamic brain abscess is a deep-seated intraparenchymal lesion that has been reported in a few of the cases in the literature.

Case description:

Herein, we present the case of a 57-year-old male patient with a two-year history of rheumatoid arthritis who was under treatment with methotrexate and etanercept for one and half years, presented to our hospital with sudden onset headache, nausea, vomiting, and fever for 5 days. Brain magnetic resonance

gT /F 1 300.63 551.67 aTJ 5T O q 0.0000093(i)6(n t)11(he U)ng (eny)4(ears)-53 O rom gin

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 081 Category: Clinical Vignette

Residency Program: Henry Ford Hospital Detroit
Presenter: Jamie Theresa Abad
Additional Authors: Katherine Heeres, Fiona Clowney MD, Eric Scher MD

A Clinical Case of Icteric Leptospirosis in an Urban Setting

Introduction: Leptospirosis, caused by the gram-negative spirochete, Leptospira, is prevalent in tropical locales. It is significantly less common in more temperate areas. Infection can present with complex clinical features ranging from a mild subclinical infection to Weil disease, characterized by multiorgan involvement, most commonly kidney and liver failure. In this report, we detail a patient admitted to a large urban hospital for management of leptospirosis, emphasizing the need for empiric therapy with strong clinical suspicion.

Case: We present a 49-year-old man with no medical history who transferred to our hospital following a 4-day stay at an outside facility. One week prior, he developed abdominal pain, fatigue, irregular bowel movements, and notable jaundice. He also reported shortness of breath, abdominal fullness, swelling of the arms and feet, and intermittent visual changes. Examination revealed jaundice, scleral icterus, bilateral lower extremity edema, and petechiae on the upper extremities. He was afebrile and hypertensive. Laboratory results revealed leukocytosis (21.6 K/ul), absolute neutrophil count (14.77), thrombocytopenia (Plt 53 K/ul), acute kidney injury (Creatinine 6.88 mg/dL) and acute liver injury with total bilirubin of 19.1 mg/dL, direct bilirubin of 16.4 mg/dL, alkaline phosphatase 81 IU/L, AST 53 IU/L, ALT 53 IU/L. Other testing including lipase, acute hepatitis panel, EBV/CMV, and autoimmune liver panel were unremarkable. CTAP revealed mild splenomegaly and MRI demonstrated hepatic steatosis without cirrhosis and two indeterminate liver lesions, likely benign hemangiomas.

Common causes of acute biliary and hepatic etiologies were ruled out. He reported insignificant alcohol use, no supplement, analgesic or illicit drug use. The patient reported a recent camping trip in Northern Michigan, involving camping, fishing, a swim in the Great Lakes, and canoeing in wooded areas along a river. He denied tick exposure or rash but mentioned a recent rat outbreak in his neighborhood. Despite no direct contact with rats, his dog killed two, and he disposed of them using a plastic bag.

Despite the absence of certain classic signs like conjunctival suffusion, the patient exhibited a constellation of symptoms and possible exposure consistent with leptospirosis. IV Doxycycline was initiated for high suspicion of icteric Leptospirosis while awaiting results, leading to significant improvement in symptoms, hyperbilirubinemia, and creatinine. Post-discharge, results returned positive for acute leptospirosis IgM, which confirmed the diagnosis.

Discussion: Early symptom recognition is crucial, especially in patients with multiorgan involvement and specific exposures. Most cases are mild, with only 10% progressing to severe forms (Weil). Leptospira transmission involves contact with contaminated soil, food, or water from urine of infected animals, predominantly rodents, entering through skin or mucous membranes. High suspicion is essential for prompt therapy to prevent further sequelae. This case emphasizes the significance of thorough history taking, particularly in non-tropical urban

Resident Poster # 082 Category: Clinical Vignette

Residency Program: Henry Ford Hospital Dro ETHQdET Q EMC /Span <</MCID 4/Lang5(en-US)>> BDC q 0.00000912 0

Resident Posters

Resident Poster

Resident Poster # 084 Category: Clinical Vignette#</MCID 1/Lang (en-US)>.9 612 792 re W* n BT /F2 1

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 087 Category: Clinical Vignette

Residency Program: Henry Ford Hospital Detroit
Presenter: Kyle

2024 Michigan Chapter ACP and SHM Michigan Chapter

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 090 Category: Clinical Vignette

Residency Program: Henry Ford Hospital Detroit
Presenter: Alyssa Naimi
Additional Authors: Elizabeth Ronchetto M.D. , Jonathon Major M.D., Zain Azzo M.D.

Posterior ST elevation Myocardial Infarction from Left Ventricular Thrombus Embolization

Background: Left ventricular thrombus is a well documented complication of patient's with reduced ejection fraction. The literature supports anticoagulation of left ventricular thrombus to reduce the risk of embolization. Cerebral emboli remain the most common complication from left ventricular thrombus, however, rare complications including coronary, splenic, renal, and limb embolism leading to acute ischemia are less commonly documented sequalae. We present a case of a young man with a posterior STEMI secondary to coronary embolism of left circumflex from left ventricular thrombus.

Case presentation: A 29 year old man with a past medical history remarkable for heart failure with improved ejection fraction presented to our institution's emergency department with chest pain, diaphoresis, and shortness of breath that had been ongoing for a 3 hours. On physical examination, he was afebrile, regular heart rate, and tachypneic. He appeared uncomfortable and diaphoretic with trace pitting edema.

Initial laboratory workup revealed a high sensitivity troponin value trend of 28, 32, and 130 which peaked at >20,000 within 10 hours of presentation. Brain natriuretic peptide was 1,567. Hemoglobin was 6 from a normal baseline. EKG demonstrated ST depression in V2V4 consistent with a posterior STEMI.

Given the laboratory and EKG findings, acute coronary syndrome was suspected. However, due to concern for bleeding with abnormal hemoglobin as well as history of normal coronary arteries on cardiac catheterization 2 years prior, the decision was made not to move forward with coronary angiogram with or without percutaneous coronary intervention. He was admitted to the intensive care unit for further monitoring. His repeated hemoglobin was within normal limits. He was started on dual anti platelet therapy and intensity heparin. Echocardiogram demonstrated ejection fraction of 8% (severely reduced), large fixed apical thrombus 2.9 cm in size, Grade III diastolic dysfunction, severely dilated left and right atrium, pulmonary artery pressure of 41 mmHg. When compared to echocardiogram 2 years prior, the ejection fraction was reduced and there was a new apical thrombus present.

Following stabilization of anemia, 3 days after presentation, patient underwent coronary angiogram which was remarkable for mid left circumflex 90% stenosis, 2nd obtuse marginal 71 % stenosis, 3rd obtuse marginal 90% stenosis, posterior left ventricular artery 99% stenosis. Findings were consistent with thrombotic embolization likely secondary to left ventricular thrombus into the distal left circumflex, posterior left ventricular artery, and 2nd obtuse marginal branch. Given the late findings and lack of current symptoms, the decision was made to treat medically with anticoagulation.

Clinical implications: Clinicians should be aware of the cardiac risks of left ventricular thrombus and keep left ventricular thrombo-embolism on the differential for patients presenting with a ST elevation myocardial infarction. In our case, a particularly high level of suspicion was warranted given patient's normal coronary arteries prior and new left ventricular thrombus on echocardiogram. Additionally, our case highlights the importance of identifying patient's with risk factors to develop left ventricular thrombus, those with reduced ejection fraction, and echocardiographic screening.

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 091 Category: Clinical Vignette

Residency Program: Henry Ford Hospital Detroit
Presenter: Elizabeth Ronchetto
Additional Authors: Jonathan Major, D.O., Alyssa tho

2024 Michigan Chapter ACP and SHM Michigan Chapter

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 094 Category: Clinical Vignette

Residency Program: Henry Ford Hospital Jackson
Presenter: Noura Attallah
Additional Authors: Iman El Feki, Tarvinder S Matharu

Staphylococcus aureus infection-related glomerulonephritis with dominant IgA and C3 deposition: case report and review of literature

Two distinct types should be considered when evaluating bacterial infections and the development of glomerulonephritis (GN): postinfectious GN, which arises weeks after infection resolution and is mostly Staphylococcus infection-mediated GN (SIAGN), and GN associated with current infection, mainly immunoglobulin A (IgA)-dominant SIAGN.

IgA primarily exists in mucous secretions and plays a critical role in establishing the microbiome, tolerating benign environmental pathogens, and locally defending against infections. One of the main causes of renal failure and glomerulonephritis is IgA nephropathy. Hematuria and Q q 0.ues of renal failure and

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 095 Category: Clinical Vignette

Residency Program: Henry Ford Hospital Jackson
Presenter: Jahnavi Ethakota
A

Resident Poster # 096 Category: Clinical Vignette

Residency Program: Henry Ford Hospital Jackson
Presenter: Parneet Kaur Hari
Additional Authors: Zarqa Yasin, MD, Jahnavi Ethakota, MD, Zreik Hassan, MD, Bipneet Singh, MD, Nadeem Ullah, MD

A Rare Case of Spontaneous Recurrent Pneumoperitoneum Attributed to Jejunal Diverticular Micro-perforation

INTRODUCTION: While most pneumoperitoneum cases arise from visceral perforations necessitating emergent surgery, approximately 5-15% result from non-surgical causes. Among these, small bowel diverticular micro-perforations are notably rare, with jejunoileal diverticulosis occurring in only 1-2% of cases posing significant diagnostic and management challenges.

CASE: We present the case of an 86-year-old female with a medical history of GERD, chronic idiopathic constipation and a surgical history of colectomy and AAA repair. She presented to the ED with acute abdominal pain and bloating without associated emesis or diarrhea. Her vital signs were stable and the examination revealed generalized abdominal tender4(l)57isend2 re significant t[(-)abdomao(-)abdott12stic i: W* n BT Jejunala8q.4 Tm 0 g 0 G

Resident Poster # 097 Category: Clinical Vignette

Residency Program: Henry Ford Hospital Jackson
Presenter: Abraham Kisule
Additional Authors: Akhil Rahman MD, Ciji Robinson MD, Niroshan Ranjan MD, Sakshi Bai MD

I

Introduction: Endophthalmitis is an eye infection that results from bacterial, viral, fungal, or parasitic infection. Most cases are exogenous and occur after penetrating eye trauma, following eye surgery, or as an extension of corneal disease. An increasing number of instances arise after intraocular injections with anti-vascular endothelial growth factor (VEGF) medications. Other infections are acquired endogenously from hematogenous seeding from another location.

Case: This abstract highlights a case of endophthalmitis diagnosed in a 35-year-old female who presented to our community hospital to be evaluated for concerns of three days of progressively worsening fatigue and weakness. Evaluation in the ED revealed that she was septic with no apparent source of infection. She was then admitted overnight for treatment with antibiotics. Throughout the night, her vision continued to deteriorate to a point that she could hardly see. The provider caring for her performed a physical exam of her eyes the following day. A physical examination of her eyes revealed erythema of bilateral conjunctiva and yellowish discharge from her eyes. Her blood cultures collected on presentation to the Emergency Department were positive for G streptococcus. Given these findings, there was increasing suspicion that the patient had endophthalmitis, and this may have been the source of the sepsis. Given our community hospital's lack of ophthalmology services, arrangements were made to transfer the patient to a tertiary care facility with in-house ophthalmology services. A diagnosis of endophthalmitis was made at the tertiary care facility. The patient received an anterior chamber tap

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 098 Category: Clinical Vignette

Residency Program: Henry Ford Hospital Jackson
Presenter: Danesh Kumar
Additional Authors: Jahnavi Ethakota, Sakshi Bai, Manesh Kumar, Aamir Ameer, Zeeshan Arshad

A rare cause of vitamin B 12 deficiency

Presentation:
A 31-year-old male with a history of polysubstance use and bipolar disorder presented with a 1-week history of ambulatory difficulty, trouble holding objects, loss of balance, and mild paresthesias in the fingers and toes. The physical examination further revealed slow fine coordinated movements (finger tapping) bilaterally, significant dysmetria on finger to nose and heel to shin, unsteady gait, asymmeytric bilateral upper and lower extremity weakness, and no achilles reflex bilaterally. Blood work showed hemoglobin of 13.3 with MCV 93.2, Vitamin B12 level <150 pg/ml (232-1245 pg/mL), homocysteine >44umol/L (3-12umol/L), and methylmalonic acid 1.22umol/L(<0.4umol/L), UDS only positive for benzodiazepines. MRI spine shows vague and symmetric T2 signal hyperintensity within the dorsal columns of the spinal cord as noted in the cervical spine that is in keeping with underlying subacute combined degeneration.

Detailed history, workup, and imaging further ruled out autoimmune disease, malabsorption, infection, trauma, vascular disease, and metabolic causes of vitamin B12 deficiency as well as any other possible synergic cause of presentation. Upon interviewing multiple times on the latter days of admission, the patient did admit that he was using the whippet pretty heavily or at least daily before presentation to the hospital. That is the inhalation of nitrous oxide.

The patient was treated with a high dose of IV vitamin B12 and was discharged from the hospital. Patient on follow-up after a year the patient recovered completely with abstinence from NO inhalation, B12 supplements, and physiotherapy.

Discussion
Subacute combined degeneration of the dorsal (posterior) and lateral columns (white matter) of the spinal cord due to demyelination is the classic neurologic presentation of vitamin B12 deficiency that commonly presents with progressive weakness, ataxia, and paresthesias. Identifying the etiologies of vitamin B12 can sometimes be challenging for an internist. Among the rare causes of vitamin B12, NO poisoning is one of the established causes. Neurologic symptoms associated with nitrous oxide use have been attributed to vitamin B12 deficiency. Through oxidation, nitrous oxide inactivates vitamin B12.

The neurological symptoms of vitamin B12 deficiency are unspecific and can be irreversible. Early detection is therefore important, using the most sensitive and specific markers available. Individuals can present with increased levels of homocysteine and methylmalonic acid in addition to low vitamin B12 levels. Classic MRI findings associated with nitrous oxide-induced vitamin B12 deficiency include T2 hyperintensity in the posterior columns, usually over several vertebral segments, with or without involvement of the lateral corticospinal tracts. These findings represent myelin sheath degeneration. Treatment should be with IV vitamin b12 supplement and response is mainly monitored with symptomatic improvement however can also be monitored with serum levels of vitamin b12, homocysteine, and methylmalonic acid levels.

Conclusion
There is no available test to screen for nitrous oxide use. Therefore it is vital to get a detailed history regarding illicit drug use in every patient and specifically about nitrous oxide use in those presenting with neurologic

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 099 Category: Research

Residency Program: Henry Ford Hospital Jackson
Presenter: Muhammad Ahmad Qureshi
Additional Authors: Luqman Munir, Ali Akram Qureshi, Kartik Gupta, Omair Ahmed, Haseeb Tareen, Affaf Shahid, Zainab Alvi, Zarqa Yasin, Richard Santos

Temporal Trends in Alcoholic Cardiomyopathy-Related Mortality Rates from 2000 to 2020: An Analysis of Epidemiology Disparities in the United States

Background: Alcoholic cardiomyopathy (ACM) is one of the major causes of mortality in the United States. This study examines ACM-related mortality (ACMRM) trends from 2000 to 2020, using age-adjusted mortality rates (AAMR) to assess disparities among key demographic and geographic subsets.

Methods: We conducted an extensive cross-sectional analysis using the Centers for Disease Control and Prevention Wide-

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 102 Category: Clinical Vignette

Residency Program: Henry Ford Hospital Macomb
Presenter: Ahmed Abdelaziz
Additional Authors: Dr. Mohamed Ahmed, Dr. Fatima Charara, Dr. Niluka Weerakoon

A rare case of coccidioidomycosis induced acute pericarditis

Coccidioidomycosis is a fungal infection caused by Coccidioides Immitis and Coccidioides Posadasii, acquired by inhalation of fungal spores in the air. It is observed in the Southwest United States specifically in California, Texas, Arizona, New Mexico, Nevada, Utah, and Central and South America. This a case of a young immunocompetent adult who initially presented with acute pericarditis and was subsequently diagnosed with Coccidioidomycosis.

A 36-year-old male patient presented to the ED for evaluation of intermittent substernal chest pain lasting 2-3 hours radiating to his back. Chest pain was worse with deep inspiration and lying flat, and improved with sitting up. The patient also had fever, chills, diaphoresis, and myalgias. The patient is a truck driver and recently returned from Texas.

Vital signs included a temperature of 101.9 F, Pulse 124 BPM, and Respiratory Rate 26. Labs were notable for leukocytosis of 14.2, Troponin peaked at 528. Electrocardiogram showed sinus tachycardia with possible pericarditis and diffuse ST segment elevation in leads I, II, aVF, V4-

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 103 Category: Clinical Vignette

Residency Program: Henry Ford Hospital Macomb
Presenter: Muhammad Ahmed
Additional Authors: Oumair Aejaz, MD

A rare case of streptococcus pyogenes meningitis in a healthy female after a lumbar epidural steroid injection

Introduction:

Streptococcus pyogenes (GAS) is an atypical and rarely reported cause of bacterial meningitis, especially in adults. Streptococcus pyogenes is well known for causing a spectrum of diseases ranging from tonsillopharyngitis to severe invasive diseases like necrotizing fasciitis and STSS, however, rarely causing bacterial meningitis. Although rare, GAS meningitis is associated with a high case fatality rate and can result in severe neurological complications. Streptococcus pyogenes is universally susceptible to all beta lactam antibiotics, with no published reports of resistance to date. Therefore, majority of individuals respond to penicillin G as a single agent. We present a rare case of Streptococcus pyogenes causing meningitis in an adult following Lumbar Epidural Steroid injection.

Case presentation:

A 66 y.o. Caucasian female with a history of arthritis, hypertension, hyperlipidemia was admitted with headache and rigors for 5 days. A week prior, patient received a Lumbar epidural steroid injection for chronic back pain. Two days later, patient developed new onset headache that she initially attributed to the injection. In next few days headache intensified and was also associated with nausea, neck pain, dizziness and photophobia.

Patient was in acute distress and toxic appearing, however alert and oriented. Patient was febrile at 103.1° F and HR 117 bpm. Physical examination revealed positive Nuchal rigidity, Brudzinski's and Kerning sign. Otherwise, no focal neurologic signs were noted. Laboratory tests showed leukocytosis of 23.9, CT head without IV contrast showed no acute intracranial hemorrhage or mass effect. Ct lumbar spine without IV Contrast was negative for any infection. Patient was empirically treated with antibiotics including ceftriaxone, vancomycin, ampicillin, acyclovir and dexamethasone. Lumbar Puncture with CSF analysis was compatible with bacterial meningitis. CSF Gram stain showed gram positive cocci in pairs and subsequently grew many streptococcus pyogenes (Group A). Blood cultures were negative. Antibiotics were de-escalated to IV Penicillin G for 14 days. Patient responded well to treatment with remission of fever, resolution of meningeal signs and symptoms, normalization of leukocyte count. Patient was discharged home to complete 14-day course of IV Penicillin G via Midline. Patient did not develop any neurological complications.

Discussion :

The incidence of acute bacterial meningitis is 1.5/100 000 persons per year in high income countries, resulting in 15 000-25 000 cases in the US annually. Most common pathogens include Streptococcus pneumoniae (72%) and Neisseria meningitidis (11%) in people older than 16 years. Escherichia coli and Streptococcus agalactiae cause about 35% of cases of early-onset neonatal meningitis. GAS accounts for less than 1% of cases of bacterial meningitis.

Resident Posters

Resident Poster # 105 Category:

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 106 Category: Clinical Vignette

Residency Program: Henry Ford Hospital Macomb
Presenter: Areej Zaheer
Additional Authors: Harini K. Venkatesh MD , Marrouf Azar, MD, Niluka Weerakoon, MD, Ashish Verma, MD, Vincent Abejuela, D.O.

Medullary thyroid carcinoma presenting with severe hypokalemia

Introduction: Medullary thyroid carcinoma (MTC) accounts for about 1-2% of all thyroid malignancies.

70 % of most medullary thyroid carcinoma are sporadic in nature. The typical age at initial presentation ranges between 41 and 55 years.

Aside from signs and symptoms of Cushing's disease, other clinical features included generalized weakness, exertional dyspnea, distal paresthesia, galactorrhea, back or hip pain and pathological fractures with metastases to the neck, mediastinal lymph node, liver, bones and adrenals.

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 107 Category: Clinical Vignette

Residency Program: Hurley Medical Center
Presenter: Sylusha Gadipudi
Additional Authors: Yashitha Chirumamilla, Philip Mc Donald.

Fatal cardiac tamponade following cardiac ablation procedure.

2024 Michigan Chapter ACP and SHM Michigan Chapter

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 109 Category: Research

Residency Program: Hurley Medical Center
Presenter: Nikky Maharjan
Additional Authors: Calvin Ghimire, Bibek Karki, Philip McDonald

Cardiac-Related death in US Adult Cancer patients: A SEER Based Study

Introduction

Heart disease and cancer rank as the top leading cause of death in the United States and globally and remain the primary barrier to increasing life expectancy. Previous studies have suggested cardiac death in cancer patients can be primarily attributed to cardiotoxicity associated with various anticancer treatments, including chemotherapy, radiotherapy, endocrine therapy, or surgery. We report cancer-specific causes of mortality attributed to heart disease in the USA between 2000-2020 in adults 20 years or older.

Methods

Surveillance, Epidemiology, 0 0 (/F2 10 Tf ID 10/Lang (en-US)>> BDC q 0.00000912 0 612 792 re W* n BT /F2 10 Tf 1

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 110 Category: Clinical Vignette

Residency Program: Hurley Medical Center
Presenter: Ajit Brar
Additional Authors: Ajit Brar, Mohammd Omer, Philip McDonald

Yasaman Navari

A case of pneumothorax secondary to iatrogenic esophageal perforation in a 96 years old male

Introduction:

Nasogastric tube has a key role in the management of hospitalized patients, particularly the critically ill. Significant complications include pneumothorax, subcutaneous emphysema, and respiratory failure. Risk factors for complications include altered mental status, absence of a gag reflex, presence of an endotracheal tube, supine position, and head and neck surgery. We will present a case of iatrogenic pneumothorax secondary to esophageal perforation

Case presentation:

A 96 year old female with a past medical history of coronary artery disease, hypertension, aortic stenosis, gastroesophageal reflux disease, prior occipital lobe cerebrovascular accident with PEG tube dependence presented to the ED with a PEG tube malfunction, surrounding skin excoriation and pus. PEG tube was successfully removed and the site was sutured under endoscopic guidance. During manual nasogastric feeding tube placement, an iatrogenic esophageal perforation was noted distal to the upper esophageal sphincter muscle that was not amenable to endoscopic closure. Subsequently the patient developed subcutaneous emphysema. Subsequent CT scan noted a large pneumomediastinum extending diffusely into the neck, chest wall, face including the orbit and evenperitoneal cavity and left pneumothorax. Trauma surgery was consulted and the patient underwent chest tube insertion with a surgical intervention planned later for perforation repair. after discussion with the family, they decided to change patient code to comfort care only and do not proceed with any intervention

Discussion:

Iatrogenic pneumothorax is a rare but potential complication of nasogastric tube insertion2. Clinicians need to be aware of this risk, especially in unconscious or obtunded patients3. Early detection and management are crucial to prevent further complications

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters
Resident Poster # 111 Category: Clinical Vignette

Residency Program: McLaren Flint
Presenter: Rabia Latif
Additional Authors: khurram arshad MBBS, William Lim MD, Yazan Alamro MBBS, Ahmad Munir MD.

A case of Platypnea-orthodeoxia syndrome secondary to PFO with reversal of symptoms with Intervention

Introduction:

Platypnea-orthodeoxia syndrome (POS) is a medical condition causing dyspnea, where patients experience breathlessness when standing or sitting, which is relieved by lying down. This is the opposite of orthopnea. POS is characterized by decreased blood oxygen saturation upon changing from a supine to an upright position.

Case Presentation

A 79-year-old female with a significant past medical history of deep Vein Thrombosis/pulmonary embolism on warfarin, a history of breast cancer status post lumpectomy and chemotherapy, hypertension, and CKD presented to the hospital from a living facility with the symptoms of persistent hypoxemia and platypnea-orthodeoxia syndrome. The results of the physical examination and laboratory tests were normal and did not reveal any notable abnormalities or concerns. An echocardiogram revealed severe patent foramen ovale (PFO) with right to left shunt.PFO was closed using the GORE CARDIOFORM septal occluder. After the procedure, the patient's oxygenation greatly improved, and her symptoms of POS disappeared. She was discharged two days later.

Discussion:

Platypnea-orthodeoxia syndrome (POS) is a rare condition characterized by difficulty breathing and low oxygen levels when standing upright. It is often underreported due to the challenge in diagnosis. POS is typically caused by the mixing of deoxygenated venous blood with arterial blood, facilitated by abnormalities in the heart or lungs. Two mechanisms, intracardiac and intrapulmonary, contribute to this condition. The intracardiac mechanism involves the transfer of deoxygenated blood between atria through defects like patent foramen ovale (PFO). Intrapulmonary mechanism occurs due to various pulmonary abnormalities. In this case, a patent foramen ovale on a bubble study and the improvement in symptoms and oxygen requirement after closing PFO indicate that platypnea-orthodeoxia syndrome was due to the PFO. It is common for around 25% of adults to have Patent Foramen Ovale (PFO). This condition happens when the foramen ovale, a small opening between the right and left atria, remains open despite the drop in pulmonary resistance and blood pressure on the right side of the heart after birth. The most well-established complication of PFO is cerebrovascular accident, but it has also been associated with other adverse neurological, embolic events, and hypoxemic symptoms such as POS.SCAI guideline suggests PFO closure in patients with POS if another explainable cause of POS is not found. In a study called "Percutaneous Intervention to Treat Platypnea-Orthodeoxia Syndrome," 52 patients were treated with percutaneous closure of an interatrial communication between January 1997 and July 2015. The study showed that patients with platypnea-orthodeoxia syndrome can be successfully treated with a percutaneous intervention, which may require different types of devices.

Clinical significance:

Platypnea-orthodeoxia is a rare condition that causes shortness of breath (dyspnea) and is often not recognized initially. This condition occurs due to a mismatch between the pulmonary and systemic blood flow, which leads to arterial desaturation in the upright position.. Awareness of this condition is crucial to prevent delays in appropriate management. Early diagnosis and treatment can effectively alleviate symptoms and prevent further complications arising from this condition.

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day

Resident Posters

Resident Poster # 112 Category: Clinical Vignette

Residency Program: McLaren Oakland
Presenter: Heba Al Sabbagh
Additional Authors: Lara Bitar, Gretchen Yormick, Mohammad Saleh

A case report of upper extremity dvt presenting as metast12(o)-aB3GretalA d4( )26(p)-12(r)5(e)3(sen)-1

Resident Poster # 114 Category: Clinical Vignette

Residency Program:

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 115 Category: Research

Residency Program: Michigan State University Sparrow Main Hospital
Presenter: Zan Siddiqi
Additional Authors: Min Choon Tan, Yong Hao Yeo, Boon Jian San, Sze Jia Ng, Mohammad Saad Salam, Karuna Rayamajhi, Zan Siddiqi, Amey Joshi, Sarah Paria Alizadeh, Wan E Kuo, Georgette Nader, Kevin Watat, Jian Liang Tan, Jasjit Walia, Addi Suleiman, Joaquim Correia

Qi Xuan Ang

Impact of Sarcoidosis on In-Hospital Outcomes among Patients with Atrial Fibrillation: a Nationwide Readmissions Database Analysis

Introduction: Sarcoidosis is a disease that involves multiple organs, particularly the cardiovascular system. While cardiac sarcoidosis has been increasingly recognized, the impact of sarcoidosis on atrial fibrillation (AF) is not well established.

Objective: This study aimed to analyze the impact of sarcoidosis on in-hospital outcomes among patients admitted for a primary diagnosis of AF.

Methods: Using the all-payer, nationally representative Nationwide Readmissions Database, our study included patients aged 18 years or older admitted for AF between 2017 and 2020. We stratified the cohort into two groups Presen

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day

Resident Posters

Resident Poster # 116 Category: Clinical Vignette

Residency Program: Michigan State University Sparrow Main Hospital

Presenter: Hanna

2024 Michigan Chapter ACP and SHM Michigan Chapter

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day

Resident Posters

Resident Poster # 119 Category: Clinical Vignette

Residency Program: Michigan State University Sparrow Main Hospital
Presenter: Jason Law
Additional Authors:

Resident Posters

Resident Poster # 120 Category: Clinical Vignette

Residency Program: Michigan State University Sparrow Main Hospital
Presenter: gabrielpanama
Additional Authors: Adolfo Martinez M.D , Rand Sabanci M.D, Kevit Watat D.O, Moiz Saeed M.D, Andy Kim M.D, Lalitsiri Atti M.D, Darian Ang M.D, Christopher Hanson M.D

Resident Posters

Resident Poster # 121 Category: Clinical Vignette

Residency Program: Michigan State University Sparrow Main Hospital
Presenter: Rand Sabanci
Additional Authors: Moiz Saeed, MD. Dina Shaban, MD. Svietlana Masoh Vincents, MD. Divij Jha, MD. Kevin Watat, DO. Andrew Kim, MD. Gabriel Panama, MD. Harith Ghnaima, MD. Supratik Rayamajhi, MD.

Mediastinal Intrusion: A Case Report on the Unusual Course of Pancreatic Pseudocyst

A pancreatic pseudocyst, a recognized complication of both acute and chronic pancreatitis, refers to a confined accumulation of fluid abundant in pancreatic enzymes. Its development is attributed to the disruption of the

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 123 Category: Clinical Vignette

Residency Program: Michigan State University Sparrow Main Hospital
Presenter: Zan Siddiqi
Additional Authors: Kevin Watat DO, Gabriel Panama MD, Rand Sabanci MD, Adolfo Martinez MD, Andrew Kim MD, Moiz Saeed MD, Qi Xuan Ang MD, Karuna Rayamajhi MD, Mohammad Saad Salam MD, Supratik Rayamajhi MD

A Rare Case of E. Coli Endocarditis

Background

Infective endocarditis due to E. Coli is a rare etiology of endocarditis with a reported incidence of less than 1% of all native valve endocarditis. Urinary tract infection is known as the most common cause of endocarditis. Generally, E. Coli has a higher mortality rate compared to HACEK group gram negative bacteria.

Case

A 83-year-old male with past medical history of Parkinson's disease, hypertension, hyperlipidemia was admitted for complaints of fever and agitation. Approximately 1 hour after dinner patient became confused, somnolent, had generalized shakiness, and was warm to touch and brought to the ED for further evaluation. Upon arrival patient was febrile, tachycardic, tachypneic, and hypertensive. His initial labs showed a lactate of 9, anion gap 17, and bicarbonate of 21. Patient denied urinary symptoms or recent instrumentation, however, on further work up, laboratory investigations were significant for positive urine leukocyte esterase and nitrites, but negative urine cultures. He was started on Zosyn with persisting fevers and concern for sepsis. He eventually was diagnosed with E. Coli bacteremia, confirmed by blood cultures, and switched from Zosyn to Ceftriaxone due to sensitivities. A transesophageal echocardiogram was done to further evaluate the cause of E. Coli bacteremia that revealed a 4 mm filamentous mobile density attached to the ventricular side of non-coronary cusp of the aortic valve confirming the diagnosis of infective endocarditis. The patient was started on a 6 week course of intravenous ceftriaxone with regular follow-up. The patient gradually improved clinically with eventual bacterial clearance. A repeat transesophageal echocardiogram after 7 weeks showed complete resolution of above

Resident Poster # 124 Category: Clinical Vignette

Residency Program: Michigan State University Sparrow Main Hospital
Presenter: Eki Wari
Additional Authors: Georgette Nader, Adarsh Jha, Sumugdha Rayamajhi, Sarah P. Alizadeh, Mohammad Saad Salam, Zan Siddiqi, Karuna Rayamajhi, Satya Rijal, Qi Xuan Ang,

Atypical Hemolytic Uremic Syndrome in a Young Adult with Complex Psychiatric History: A Diagnostic Challenge

Atypical hemolytic uremic syndrome (aHUS) is a rare, life-threatening condition characterized by the triad of microangiopathic hemolytic anemia, thrombocytopenia, and acute kidney injury, often without a preceding significant diarrheal illness. Its rarity and the broad differential diagnosis make aHUS a formidable challenge, particularly in patients with complex medical histories. We report a case of aHUS in a 26-year-old male, underscoring the diagnostic challenges and the critical role of early intervention.

A 26-year-old male with a complex history of bipolar disorder, autism, Attention-Deficit/Hyperactivity Disorder (ADHD), oppositional defiant disorder, and pervasive developmental disorder presented with symptoms of perianal bleeding, itching, and significant lethargy. Initial laboratory results were within normal limits, masking the imminent diagnosis of aHUS. However, within 5 days, a marked deterioration in his condition was observed. This was highlighted by a significant drop in platelet count to as low as 22, indicating severe thrombocytopenia, alongside an acute kidney injury with creatinine levels peaking at 5.52. Hemolysis was evident from an elevated lactate dehydrogenase level of 2248 and a critically low haptoglobin level of <6. Further complicating the clinical picture, the patient's complement system exhibited substantial dysregulation: Protein S levels were alarmingly low at 15 (normal range: 63-140), Protein C levels were decreased to 30 (normal range: 70-173), total complement activity (CH50) was below 17 (normal: >41), C3 levels were reduced to 34 (normal range: 79-152), and C4 levels were undetectably low at less than 4 (normal range: 16-38), aligning with the diagnosis of aHUS. The presence of Shiga toxin-producing E. coli (STEC) introduced an additional layer of diagnostic complexity, blurring the lines between typical STEC-HUS and aHUS.

This case highlights the diagnostic complexities encountered in patients with aHUS, especially when psychiatric comorbidities obscure the clinical presentation. The absence of diarrheal prodrome and the patient's unique psychiatric profile necessitated a comprehensive diagnostic approach, ultimately leading to the diagnosis of aHUS

Resident Poster #: 125 Category: Clinical Vignette

Residency Program: Michigan State University Sparrow Main Hospital
Presenter: Kevin Watat
Additional Authors: Georgette Nader MD, Majid Yavari MD, Rand Sabanci MD, Gabriel Panama MD, Moiz Syed MD, Dina Shaban MD, Andrew Kim MD, Matthew Wilcox DO, Supratik Rayamajhi MD

Unusual Case of Cardiac Myxoma Mimicking Infective Endocarditis

Introduction:

Myxomas are the most common primary cardiac tumor with overall low incidence that usually arises from the interatrial septum and rarely from the mitral valve (MV). The clinical presentation of myxomas shares common systemic signs and symptoms with infective endocarditis (IE), rendering the latter a chief consideration in the differential diagnosis.

Case Presentation:

A 75-year-old female patient presented with a one-month history of fever, chest pain, and dyspnea. She had hypotension, fever, and elevated inflammatory markers. Electrocardiography revealed ST elevation myocardial infarction and atrial fibrillation with rapid ventricular response, necessitating an emergent but successful synchronized cardioversion. Coronary angiography showed a thrombus in the left anterior descending (LAD) artery. To preserve LAD integrity, no percutaneous intervention was performed. Transthoracic/esophageal echocardiography (TTE/TEE) revealed a large, irregular, mobile MV mass (2.7x2.3 cm) suspected, and intravenous (IV) antibiotics were initiated without improvement in infectious markers, and cultures remained consistently negative.

consistis EMC /Span <</MCID 8/Lang (en-US)>> BDC q 0.00000912 0 612 792 re W* n BT /F2 10 562(o(sm-2(s /F2 10 G.02)>> B 160

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 127 Category: Clinical Vignette

Residency Program: Trinity Health Grand Rapids
Presenter: Taylor Beckwith
Additional Authors: Ramsha Aqeel, Mark Spoolstra

An uncommon cause of B12 deficiency

Introduction: Nitrous oxide use is not often thought of as a cause of B12 deficiency. Diagnosis can be challenging in such patients and awareness is imperative for patients to receive appropriate treatment.

Case: A 43yo male with a history of polysubstance abuse, depression, ADHD, and possible seizure, presents with altered mental status. The patient was using nitrous oxide when he started to feel confused and called an ambulance. Over the past month, he has been 'sniffing' nitrous oxide canisters with increasing frequency. He also endorses having double vision and difficulty with balance. On exam, he is oriented x3, but has difficulty answering questions due to significant anxiety. He also has hyperreflexia in the bilateral upper and lower extremities and is intermittently tremulous. B12 level was found to be 113. Methylmalonic acid level was 3.24 and Homocysteine was >130. MRI brain and MRI spine were unremarkable, with no evidence of a demyelinating process. EEG showed no epileptiform abnormalities. The patient was treated with IM B12 injections. He did show some improvement in his mentation, but still endorsed difficulty concentrating and word finding difficulty. Vitamin B12 levels normalized and the patient was discharged to inpatient rehab.

Discussion: Nitrous oxide use is a lesser known cause of B12 deficiency. eA-8(>1N6(esP7g1en h41 q 0.00000912 0 612

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day

Resident Posters

Resident Poster # 128 Category: Continuous Quality Improvement/Evidence-based Medicine

Residency Program: Trinity Health Grand Rapids

Presenter: Brian Cutler

Additional Authors: Adam Anderson, Taylor Barthels, Reid Mitchell, Olawale Olanisa

Quality Review of False-Positive Cardiac

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 129 Category: Clinical Vignette

Residency Program: Trinity Health Grand Rapids
Presenter: Courtney Herron
Additional Authors: John Westfall, DO; Andrew Jameson, MD

Green Neutrophilic Inclusions: Really the "Green Crystals of Death"?

There are many types of inclusions that can be seen on blood smears that can aid in diagnosing a patient. Bright green neutrophilic inclusions are findings on a peripheral blood smear that pathologists deem a rare find but can indicate the rapid demise of a patient.

A 70-year-old female presented to the ED with recurrent abdominal pain. She was hospitalized 1 month prior to admission with a newly diagnosed pancreatic mass with good clinical response to biliary stenting via ERCP. Endoscopic biopsy was consistent with malignancy but not diagnostic for the specific type of cancer. Before she could be seen by Oncologic Surgery, she presented back to the hospital with pain, elevated alkaline phosphatase, AST, and ALT. She underwent repeat ERCP with stent exchange with clinical and laboratory improvement. Two days after the ERCP, she developed worsening pain, hypotension, and lactic acidosis requiring vasopressor support. She experienced progressive decline in mental status, acute kidney injury, and thrombocytopenia (dropping from 129 to 17 over 4 days). Due to worsening coagulation studies (PT/INR and aPTT) a manual differential was performed revealing normochromic and normocytic erythrocytes with mild anisocytosis with rare nucleated red blood cells. No schistocytes or rouleaux changes identified. Platelets were markedly reduced. Leukocytes were increased with mature neutrophils showing toxic granulation, vacuolization, and Dohle bodies. There were rare bright green-blue inclusions seen. Antimicrobials were expanded from ceftriaxone to pipercillin/tazobactam due to desire to broaden intra-abdominal coverage. The Infectious Diseases team was contacted regarding the blue-green inclusions implying very significant poor prognosis. The arrival of these inclusions has a strong indication that the patient will expire within 48 hours of the result. The patient continued to rapidly decline, and she passed away on day 6 of admission. Post-mortem the patient had blood cultures turn positive for Enterococcus faecalis and a next-generation sequencing of whole blood revealed Escherichia coli.

This further illustrates the prognostic accuracy of the appearance of green inclusions within neutrophils. This finding has been described previously in multiorgan failure and septic shock. This has been most frequently described with Escherichia coli. This case additionally shows the value of a manual differential in diagnosis of systemic illness.

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 130 Category: Clinical Vignette

Residency Program: Trinity Health Grand Rapids
Presenter: Olawale Olanisa
Additional Authors: Abiy Nigatu, MD, FACC, Abdul Rahman Omer, MD, FACC, Andre Hollingsworth, MD

Fulminant Endocarditis in a Ir0goe(P)7(o)-5en-US

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster #: 131 Category: Clinical Vignette

Residency Program: Trinity Health Grand Rapids
Presenter: Jasmine Zahid
Additional Authors: Ronald M Hofmann MD, Rehan Ansari MD

The Great Masquerader: 3 Cases of Bartonella in Renal Transplant Patients

Introduction

Bartonella infections present with a wide array of non-specific clinical manifestations including lymphadenopathy, fever, diarrhea, and hepatosplenomegaly. Literature regarding Bartonella infections in renal transplant recipients is limited. The following 3 cases demonstrate Bartonella infections following renal transplantation at a single center. Standard induction immunosuppression included Thymoglobulin and Solumedrol in all 3 patients.

Case 1

25-year-old female with a history of neonatal left renal vein thrombosis and secondary hypertensive nephrosclerosis leading to ESRD underwent a deceased donor kidney transplant (KDPI 17%) with 1 HLA-A antigen mismatch. Recipient cPRA 99% . CMV: R+. EBV: D+/R+) with an uneventful course until 1 year later. Maintenance immunosuppression was MMF, Tacrolimus and prednisone. She presented with fevers, diarrhea, generalized fatigue and an unproductive cough who was found to haven an AKI. Patient had adopted feral kittens 4 months prior to admission. She reports both kitten and flea bites. Creatinine on admission was 1.5, peaked at 1.7 and returned to her baseline of 1.1-1.3. CT showed numerous ill-defined hypo-attenuating liver lesions. MRI showed diffuse nodular lesions throughout the liver with diffuse enhancement. Concern for PTLD was high. Karius was positive for both EBV and Bartonella. Bartonella Henselae IgG antibody 1: 128 henselae IgM antibody 1:256. She was treated with doxycycline 100 mg twice da

Resident Posters

Resident Poster # 132 Category: Clinical Vignette

2024 Michigan Chapter ACP and SHM Michigan Chapter President and Medical Student Day
Resident Posters

Resident Poster # 133 Category: Clinical Vignette

Residency Program: Trinity Health Livonia
Presenter: Bazigh Naveed

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 134 Category: Research

Residency Program: Trinity Health Livonia
Presenter: Mutjaba Abdellatif
Additional Authors: Mujtaba Abdellatif, Ayman Alsaadi, Juma Rashid Bin Firos, Omar Alkhateeb, Sameer Mallick, Paul Nona, Tanmay Swadia, Preeti Misra

Ibrahim El-Mair

The Comparison of Using Single Versus Two Catheter Concept in Diagnostic Transradial Coronary Angiography: A Single Community Center Experience

Background: Coronary artery anomalies (CAAs) are congenital conditions characterized by abnormal origin or course of the 3 main coronary arteries. Most variations are benign; however, some may lead to myocardial ischemia and/or sudden cardiac arrest depending on where the artery travels.

Case: 66 year old G [(C)4(a)-5(th)-15(e)3(ter)5(C)5(o)13(n)-12(c)-6(e)3(p)12(t i)-5(n)-1wo

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 137 Category: Clinical Vignette

Residency Program: Trinity Health Livonia
Presenter: Bakht-Awar Khan
Additional Authors: Bazigh Naveed, Alea Mohamed, Hemica Hasan, Sarmad Pirzada

Not every confusion is UTI

Introduction: Invasive Listeriosis is a treatable but potentially fatal disease if not recognized early. It is a relatively rare condition that mainly affects elderly patients and immunocompromised hosts. Here we present a case of missed Listeria bacteremia in a patient with polymyalgia rheumatica on chronic immunosuppressive therapy.

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 138 Category: Clinical Vignette

Residency Program: Trinity Health Livonia
Presenter: Brock Malatches
Additional Authors: Jemeera Jeyamuhunthan MD, Sarmad Pirzada MBBS, Stefan Odabasic MD, Ayman Alsaadi MD

IVIG-Induced Aseptic Meningitis in an Adult Patient with Acute Inflammatory Demyelinating Polyneuropathy: A Case Report

Intravenous immunoglobulins (IVIG) therapy is used to treat various autoimmune, immunodeficiency, and inflammatory conditions. One of the rare, but serious, side effects is aseptic meningitis. In this case report, we present a 55-year-old female who experienced IVIG-induced aseptic meningitis for treatment of acute inflammatory demyelinating polyneuropathy (AIDP).

We present a 55-year-old female with multiple comorbidities who presented to the emergency department with a chief complaint of acute bilateral lower extremity weakness for 2 days associated with ascending paresthesia, tingling, and worsening chronic back pain. Additionally, she reported sore throat and intermittent fever in the last few weeks prior to presentation, however, she denied headache and neck stiffness. Physical examination showed tenderness overlying the thoracic spine, 4/5 strength in the bilateral lower extremities, but intact sensation.

Initial work up consisted of a comprehensive metabolic panel, complete blood count with differential, and viral panel including COVID-19 and Influenza testing, which all returned within normal limits. The patient underwent MRI with and without contrast of the entire spine which was within normal limits. A lumbar puncture (LP) was performed and was within normal limits as well. The patient was suspected to have AIDP for which she received treatment with IVIG. On day 4 of treatment, the patient was noted to have headache, neck stiffness, and chills, therefore the IVIG was stopped. A repeat lumbar puncture was performed revealing 129/mm3 total nucleated cells with 77% lymphocytes, 56/mm3 RBCs, glucose 52mg/dl, and protein 35mg/dl. Pathology review revealed small mature lymphocytes. An extensive infectious workup was performed including CSF PCR for Escherichia coli K1, H. influenzae, L. monocytogenes, N. meningitidis, S. agalactiae, S. pneumoniae, CMV, Enterovirus, HSV 1 and 2, HHV 6, VZV, and C. neoformans, which all returned negative. In addition, Gram stain and culture, and Lyme serologies were negative. With the return of the above findings, a diagnosis of aseptic meningitis induced by IVIG was made with the collaboration of a multidisciplinary team which included neurology and infectious diseases. In addition to IVIG discontinuation, supportive care was started with adequate hydration and symptomatic management with improvement of her symptoms over the following days and she was subsequently discharged to a rehabilitation facility.

IVIG-induced aseptic meningitis is a known, but relatively rare, adverse effect that must be included in the differential if a patient develops symptoms consistent with meningitis during or after treatment. The clinical presentation of aseptic meningitis is very similar to other types of meningitis and work up should include lumbar puncture to rule out infectious etiology first and foremost. CSF analysis generally shows a neutrophil predominance however this is not sensitive or specific. Symptoms usually occur within the first 48 hours after initiation of treatment, however in our case symptoms developed after 96 hours. Therefore, clinicians should have a high index of suspicion for such a complication throughout the treatment course as identification may warrant immediate discontinuation of IVIG treatment. In most cases, the symptoms are self-limited with conservative treatment, however complications may arise if not detected early.

Resident Poster # 140 Category: Research

Residency Program: Trinity Health Oakland
Presenter: Fathima Shehnaz Ayoobkhan
Additional Authors: Geetha Krishnamoorthy Departmentl

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 141 Category: Research

Residency Program: Trinity Health Oakland
Presenter: Sharanya Shree Ezhilarasan Santhi
Additional Authors: Roopali Dahiya, Aiman Perween Afsar, Amna Hashmi, Ahmedyar Hasan, Zara Arshad, Faisal Nawaz, Rahul Kashyap

Exploring The Role of Infectious Triggers in Causation of Sarcoidosis

INTRODUCTION: Sarcoidosis has complex pathogenesis, with multiple innate and external risk factors. Microbial organisms have been implicated as a trigger in those with genetic predisposition. This systematic review of case reports/series aims to determine patterns in pulmonary sarcoidosis which was diagnosed along with or post an infectious episode.

METHOD: Case reports/series published from January 2015 to September 2023 were searched through PubMed. . Subsequently, articles were exported to Rayyan software. Each study was reviewed by two investigators based on inclusion and exclusion criteria. Data from these selected case studies of patients with pulmonary sarcoidosis diagnosed alongside or after an infection, were then collected and analyzed for clinical, laboratory and radiological abnormalities. The study was IRB exempt.

RESULT: Total 21 patients were included from 19 case reports and one case series. Mean age of the patients was 45 years, ranging from 16-75 years. Total 12 of them (57 %) were male. Most common presenting symptoms were cough (47.6%) and dyspnea (43%). Comorbidities noted in the patients are Type II Diabetes Mellitus, Hypertension and Chronic Kidney disease (14% each). Stage III sarcoidosis was diagnosed in 33.3%, Stage II and concomitant extra-pulmonary involvement were seen in 28.5% each. The predominant infectious etiology recognized were Mycobacterium tuberculosis (38%), SARS-CoV-2 virus (24%) and Propionibacterium acnes (14%). 8 of 21 patients (38%) had abnormalities on chest radiography. On Computerized tomography of the chest, most common radiological abnormalities noted were nodular opacities (47.6%) and mediastinal lymphadenopathy (38%). ACE levels were abnormally elevated in seven patients and only one had hypercalcemia. Patients were treated with systemic steroids (61.9%) and methotrexate(14.2%), with clinical and radiological resolution noted in 52.3% of the study population.

CONCLUSION: Our systematic review focussed on clinical presentation, blood and radiological patterns, treatment administered in those who were diagnosed with sarcoidosis precipitated by an infection. It is important to be aware of such a complication in those who do not improve after appropriate antibiotic therapy for a recognized infectious etiology.

Resident Poster #: 142 Category: Clinical Vignette

Residency Program: Trinity Health Oakland
Presenter: Prami Nakarmi
Additional Authors: Nikhil Vojjala, MD; Sumit Raut, MD; Pritha Chitagi, MD; Geetha Krishnamoorthy, MD, FACP

Concomitant Herpes Simplex Virus (HSV) 1 and Adenovirus ARDS in an immunocompetent patient: Innocent bystander or Vicious culprit

Introduction:

Acute respiratory distress syndrome (ARDS) is associated with a very high mortality (40 to 60%). Identification of the etiology and specific treatment, in addition to guideline-based ventilation strategies improves outcomes. We present an immunocompetent woman with Herpes Simplex Virus 1 (HSV1) and Adenovirus related severe ARDS successfully managed with mechanical ventilation and early steroid therapy.

Case Presentation:

A 40-

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 143 Category: Clinical Vignette

Residency Program: Trinity Health Oakland
Presenter: Rishab Prabhu
Additional Authors: Nikhil Vojjala, MD; Sangeetha Krishnamoorthy, MD; Ibrahim Azar, MD; Muhammad Nadeem Khan, MD; Shireesh Gadgeel, MD; Geetha Krishnamoorthy, MD FACP

Reno-Duodenal fistula: An unusual cause of polymicrobial sepsis in a lung cancer patient (Turning Tumor titbits into tumor trouble)

Introduction:

Sepsis in patients with underlying malignancies can be due to many causes such as infections due to neutropenia, pneumonia, and gastrointestinal infections. Fistula formation leading to sepsis is very rare. Factors associated with gastrointestinal fistulization as a possible cause of sepsis include recent radiation therapy, gastrointestinal surgery, and polymicrobial growth on blood cultures, especially gut microbiota.

Case description:

A 60-year-old man with hypertension, Hyperlipidemia, and non-small cell lung cancer presented to the emergency department (ED) with a history of fever with chills of one day duration. Family members reported that the patient was dull appearing and had difficulty in word finding, for the past week. Of note, the patient's non-small cell lung cancer initially responded to therapy, then became resistant to multiple lines of therapy including progression while on a clinical trial. The patient was then placed on Gemcitabine-based chemotherapy and underwent radiation therapy for right renal metastasis three months before presentation as a palliative intent. On presentation to the ED, he was febrile (103.6 F), hypotensive (blood pressure 70/50 mm Hg), tachycardic (121/min), and hypoxic saturating 87% on room air. Physical examination revealed an ill-appearing man with tachypnoea. Neurological examination showed that he was oriented only to a person. There was right costovertebral angle tenderness with a resonant percussion note in the lower posterior thorax. He was resuscitated with intravenous fluids, and oxygen, and started on antibiotics as per sepsis protocol. A computerized tomography (CT) scan of the chest with an angiogram showed multiple nodules with cavitation and no pulmonary embolism. The patient's vital signs improved with fluid resuscitation and antibiotics initially. Blood cultures grew multiple atypical organisms both bacteria and fungi. Bacteria were Lactobacillus acidophilus, Streptococcus, and Leuconostoc mesenteroides, and fungi were Saccharomyces Cerevisiae and Candida Albicans. Given polymicrobial sepsis, the gut was suspected as the source. CT scan with contrast of the abdomen showed a right upper pole renal mass. Air was seen within the right renal parenchyma, with a fistula between the 1st part of the duodenum and the anterior surface of the kidney. Esophagogastroduodenoscopy confirmed the large fistulous opening in the duodenum and extending into the kidney. Endoscopic closure of the fistulous opening was attempted but was not possible due to adherence to the kidney. Transesophageal echocardiogram did not show any vegetation. A CT scan brain showed metastatic disease. Because of multiple medical comorbidities, poor performance status, and progressive disease, the patient and his family members requested hospice care.

Reno-duodenal fistula is a very rare and unusual cause of sepsis in patients with malignancy. Polymicrobial blood cultures prompted us to think about gut translocation. Early identification may make surgical intervention possible. Though in this patient, we were not able to control the source of sepsis, as a minimally invasive procedure, endoscopy should be done with an attempt to close the fistula, if possible, in poor surgical candidates. Poor performance status, diffuse metastatic disease, and recurrent hemodynamic instability excluded the surgical treatment options for this patient.

2024 Michigan Chapter ACP and SHM Michigan Chapter

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 145 Category: Clinical Vignette

Residency Program: Trinity Health Oakland
Presenter: Nikhil Vojjala
Additional Authors:

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters
Resident Poster # 146 Category: Clinical Vignette

Residency Program: Trinity Health Oakland
Presenter: Fizza Zulfiqar
Additional Authors: Fizza Zulfiqar MD, Mahim Mahim, Nikhil Vojalla MD, Geetha Krishnamoorthy MD, FACP, Pratik Bhattacharya, MD

Hyponatremia with progressive neuropsychiatric symptoms prompts investigation for LGI1 (Leucine-Rich glioma inactivated-1) antibody encephalitis?

Introduction:

Limbic encephalitis (LE), a rare neurological disorder characterized by limbic system inflammation, presents with a wide range of symptoms, including subacute cognitive decline, short-term memory loss, faciobrachial dystonic seizures (FBDS) and behavioral abnormalities (1). Etiologies include infections, tumors, and autoimmune reactions. During the disorder, 60-80% of patients develop hyponatremia (1) that might delay the diagnosis of encephalitis. The etiology for hyponatremia remains unclear but is presumed secondary to SIADH. Herein, we report a man with progressively worsening dementia and hyponatremia due to autoimmune Leucine-Rich glioma inactivated 1 (LGI-1) encephalitis.

Case presentation:

A 67-year-old man with prior stroke, hypertension, coronary heart disease, and depression presented to the emergency department due to seizure-like activity. Patient was sleeping while he experienced sudden stiffness and jerking movements of both upper and lower limbs lasting for a minute. The patient appeared confused and was brought to the hospital.

For the past nine months patient has been experiencing multiple neuropsychiatric symptoms including vivid dreams, hallucinations, delusions, mild cognitive impairment with some gait imbalance and urinary incontinence. He had a prior admission due to new-onset cognitive impairment. He was seen by neurology, and computerized tomography (CT) scan head, magnetic resonance imaging (MRI) brain and electroencephalogram were negative. Considering encephalitis, an autoimmune panel was ordered that was later positive for Leucine-Rich glioma inactivated 1 antibody (LGI-1 IgG) . On this admission with worsening mentation and recurrent seizures episodes, neurology was reconsulted.

Laboratory investigations were significant for hyponatremia (130 mmol/L), serum osmolality was 264 Mosmol/K. Further workup showed urine sodium of 49 meq/L, urine osmolarity of 470 mosmol/K and uric acid 2.8 mg/dL. Thyroid Stimulating Hormone and cortisol levels were normal. Due to euvolemic hypoosmolar hyponatremia, syndrome of inappropriate antidiuretic hormone (SIADH) was diagnosed. CT chest and abdomen was negative, Positron Emission Tomography scan of the body was normal, done in a search for primary malignancies. Repeat brain imaging was normal. SIADH was attributed to ongoing underlying LGI-1 encephalitis. The patient was started on levetiracetam 500 mg twice daily for seizure control with salt tablets and IV immunoglobulins (IVIG) for 5 days. Therewas improvement in psychiatric/behavioral symptoms and sodium trended up with fluid restriction and salt tablets. Post IVIG, the patient was started on tapering dose of oral steroid to reduce the risk of recurrent seizures, hyponatremia, and behavioral problems with advice to follow-up as outpatient with neurology.

Discussion:

LE has been considered a paraneoplastic phenomenon, most commonly in association with underlying lung cancers [2]. Recently it has become apparent that LE may occur in the absence of malignancy due to autoimmune etiology. In animal studies, LGI-1 is highly expressed in the hypothalamus, indicating a possible inflammatory response. ADH (antidiuretic hormone) secretion can be disrupted by this inflammatory reaction, resulting in

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 147 Category: Clinical Vignette

Residency Program: University of Michigan Med-Peds
Presenter: Katerina Castillo
Additional Authors: Arielle Schaffer-White, Rebecca Northway

Shoulder effusion: not a crystal clear diagnosis

Introduction:

Shoulder complaints are common in the primary care office. Pseudogout is an overlooked cause of acute monoarthritis of the shoulder. Presentation can be similar to other diagnoses, including septic arthritis, which can coexist with pseudogout.

Case Description:

An 83 year old male with a history of gout presented to the primary care clinic with acute onset right shoulder

Resident Poster # 148 Category: Clinical Vignette

Residency Program: University of Michigan Med-Peds
Presenter: Micaela Witte
Additional Authors: Dr. Michael Joynt MD

Introduction: Platypnea-orthodeoxia syndrome (POS) is a unique clinical syndrome seen in specific types of hypoxia, categorized by the decrease in oxygen saturation by more than 5% or the decrease in arterial oxygen pressure by more than 4 mmHg when transitioning from recumbency to standing. The differential is limited to intracardiac or intrapulmonary shunts. Prompt identification is crucial as delayed closure can result in progressive hypoxia, heart failure, or neurologic complications.

Case presentation: A 68-year-old woman with a notable history of hypothyroidism, 15-pack-year smoking history, asbestos exposure, and recent diagnosis of presumed provoked deep vein thrombosis (DVT)/pulmonary embolism (PE) on apixaban, presented to pulmonology for a second opinion of a three month history of persistent and progressive hypoxia requiring 4-12 L nasal cannula at home. She had previously had an echocardiogram and a CT chest at an outside hospital that were reportedly unremarkable. During her pulmonary function testing (PFT), which was remarkable only for persistent hypoxia, she was noted to require up to 15L nasal cannula to maintain saturations >90%, so she was sent to the emergency department for further evaluation. Initial evaluation, including VBG, CBC, CMP, TSH, ESR, BNP, and troponins as well as EKG were unremarkable. CTPE showed scattered bronchial thickening in the lung bases, but no evidence of clot.

The patient was admitted for further evaluation at which point she was noted to have worsening hypoxia and increased oxygen requirements when sitting or standing, consistent with POS syndrome. Further evaluation with high-resolution CT scan showed mild interstitial lung disease and deep venous ultrasound was negative for residual DVT. Notably, transthoracic echocardiogram revealed a large right to left shunt on bubble study. Cardiology was consulted, at which time cardiac MRI was ordered and showed a large sinus venosus atrial septal defect (ASD) with

Resident Posters

Resident Poster # 149 Category: Clinical Vignette

Residency Program:

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 150 Category: Clinical Vignette

Residency Program: Wayne State University Detroit Medical Center Sinai Grace
Presenter: Nizar Alnabahneh
Additional Authors: Muslehuddin, Zainab MD; Raied, Hanna MD

Glomerulopathy in the setting of Morbid Obesity with controlled Diabetes

Introduction: Obesity is a very common and complex presentation in the US. Class 3 obesity is defined as a BMI of 40 or higher. The prevalence of obesity is rapidly increasing across the world and all sections of society. Obesity related renal disorders have surged tenfold in recent years, with obesity causing increased glomerular filtration rate (GFR) and glomerulomegaly. In type 2 diabetes, heightened hyperfiltration leads to irreversible kidney damage and eases the progression to end-stage renal disease (ESRD). The initial presentation is nephrotic range proteinuria and can be distinguished from Focal segmental Glomerulosclerosis (FSGS) by glomerulomegaly and less severe foot process effacement band without a complete nephrotic syndrome. Further studies are needed to explain why patients with obesity-related glomerulopathy do not develop complete nephrotic syndrome.

Case Description: A 48-year-old male with a past medical history of type 2 diabetes mellitus, obesity class III, and associated chronic kidney disease stage 3, presented to the emergency department complaining of increased lower extremities bilateral swelling, exertional dyspnea, and reduced urine output over the last week. Upon examination, he exhibited a significantly marked bilateral pitting edema in the lower extremities. Vital signs on admission were within normal limits. However, his BMI measured 54.6

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 151 Category: Research

Residency Program: Wayne State University Detroit Medical Center Sinai Grace
Presenter: Ajmat Ansari
Additional Authors: Ajmat Ansari MD, Malitha Hettiarachchi MD, FACP

Comparative Efficacy of Spironolactone and Eplerenone in Chronic Heart Failure

Introduction: Chronic heart failure is a prevalent clinical syndrome associated with significant morbidity and mortality, affecting approximately 6.5 million Americans over the age of 20. Mineralocorticoid receptor antagonists (MRAs) have been integral to HF therapy for the past 16 years, with spironolactone and eplerenone being the recommended options. This systematic review aims to contribute evidence-based insights into the comparative efficacy of these MRAs specifically Spironolactone and Eplerenone, in the treatment of chronic heart failure (HF). The primary endpoint evaluated is all-cause mortality, with secondary endpoints including left ventricular ejection fraction (LVEF) improvement, cardiovascular mortality, and HF-related hospitalization.

Methods: A comprehensive search strategy was employed, utilizing PubMed to identify randomized controlled trials (RCTs) conducted from 2010 to 2023. The inclusion crit2usignificant morbidity and

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 152 Category: Clinical Vignette

Residency Program: Wayne State University Detroit Medical Center Sinai Grace
Presenter: Pradeep Balasubramanian
Additional Authors: Kausar Hafeez MD; Wasif Hafeez MD

Shagreen patch on the face in a patient with Tuberous sclerosis Complex - An Uncommon Site

Introduction:

Tuberous Sclerosis Complex (TSC) is a rare neurocutaneous disorder that includes multiple variations of skin changes that can occur throughout the body. The most common skin findings include Ash leaf marks which are often present from birth; facial angiofibroma's which commonly occur around the cheeks and nose starting in childhood, cephalic plaques on the forehead that are raised and fibrous, Shagreen patches which are commonly found on the torso or lower back, and ungual fibromas around or under the nails. While these skin conditions can commonly be found in patients with TSC, it is rare to find Shagreen patches outside of the torso or lower back. Here, we present a case of a Shagreen patch found on the patient's face.

Case Report:

A 53-year-

Resident Posters

Resident Poster # 155 Category: Clinical Vignette

Residency Program: Wayne State University Detroit Medical Center Sinai Grace
Presenter: Suhitha Bysani
Additional Authors: Kiani, A.; Solanki, N.; Chaudhary, A.; Bilal, M.

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 156 Category: Clinical Vignette

Residency Program: Wayne State University Detroit Medical Center Sinai Grace
Presenter: Howard Dabbous
Additional Authors: Raza Ur Rahman, Mohammad Azfar Bilal, Mohamed Siddique

Obscure Gastrointestinal Bleeding in a middle age man

Introduction: The most common type of gastrointestinal bleed (GIB) is an upper GIB. Incidence of UGIB is approximately twice that of lower and presents with hematemesis and melena while lower GIB commonly presents with hematochezia. Previously, approximately 5-10% of patients were labeled as obscure bleed after no source of bleeding being found on standard endoscopic and radiology evaluation. However, in 75% of these patients, the source of bleeding is the small intestine that is not seen on standard evaluation. Advancements in technology have changed the definition of obscure bleed to comprise only patients with no identifiable bleed source after complete small bowel evaluation in addition to standard endoscopic and radiographic evaluations.

Case Description: A 59-year-old male presented to the emergency department (ED) after an episode of massive hematochezia and syncope. Patient has no history of prior GIB and did not report dizziness, dyspnea, or fatigue. Patient's vital signs were initially stable with laboratory studies showing hemoglobin (Hb) of 9.1 g/dl. Endoscopy demonstrated a small arteriovenous malformation with no active bleeding and colonoscopy showed dark stool without active bleed. Patient continued to have hematochezia with a nadir Hb of 4.6 g/dl and received a total of 5 units of packed red blood cells (PRBCs). CT angiography (CTA) abdomen/pelvis, radionuclide GI bleed scan, and a gastrografin enema were all negative for any pathology or bleed. Hematochezia resolved and the patient was discharged with a Hb of 7.6 g/dl. Patient returned to the ED 2 days afterwards complaining of lethargy, dizziness, weakness, and melena. Patient was hypotensive on presentation (79/54 mmHg) with HR of 84bpm and digital rectal exam revealing black stool. Hb was 3.7 g/dl and the patient was admitted directly to the MICU. CTA abdomen/pelvis showed no active bleed and invasive angiograms of the celiac, superior mesenteric, and inferior mesenteric artery and their main branches revealed no active bleed as well. During the second admission, patient received 6 units of PRBCs with melena resolving, vitals stabilizing and the patient was eventually discharged. Patient was referred to a tertiary care center if symptoms returned. 5 days later, patient presented to an outside hospital with same symptoms of melena and syncope. Patient was resuscitated with 1unit of PRBC and received capsule endoscopy that showed no active bleeding. Patient was discharged home with GIB etiology remaining unknown.

Discussion: This case demonstrates elusiveness in diagnosing etiology of GIB when it is not localized to the stomach or colon. Physicians need to consider transferring the patient to a tertiary care center for further evaluating directly to the MICU.

Resident Poster # 158 Category: Clinical Vignette

Residency Program: Wayne State University Detroit Medical Center Sinai Grace
Presenter: Naisargee Solanki
Additional Authors: Naisargee Solanki MD, Zainab Muslehuddin, MD, Montaser Elkholy

Resident Poster #

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 160 Category: Clinical Vignette

Residency Program: Wayne State University Detroit Medical Center Sinai Grace
Presenter: Manisha Guntha
Additional Authors: Fred H. D. Charles MD MPH; Malitha Hettiarachchi, MD FACP

Lower extremity Hemiparesis - It is not a pinched nerve but Neurosarcoidosis

INTRODUCTION

Sarcoidosis is a multi-organ granulomatous disease of unknown etiology. It mostly affects middle-aged patients, and prevalence is higher in women compared to men. Neurosarcoidosis is relatively uncommon with a reported prevalence of 3-10%. Any part of the nervous system can be affected including brain parenchyma, cranial nerves,

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 161 Category: Continuous Quality Improvement/Evidence-based Medicine

Residency Program: Wayne State University Detroit Medical Center Sinai Grace
Presenter: Zohaib Khan
Additional Authors: Fred H. Charles, Malitha Hettiarachchi

Heart failure outcomes in GDMT alone versus GDMT combined with revascularization in patients with HFrEF

Background:

Worldwide, 5.7 million individuals are affected by heart failure, with an expected increase to 8 million by 2030. Currently, guidelines dictate that all patients with heart failure be started on guideline-directed medical management (GDMT). However, given that the most common cause of heart failure remains ischemic, this study aims to assess the benefits of adding revascularization as a treatment option for patients with HFrEF who are already on GDMT.

Methods:

A comprehensive literature search using databases such as PubMed and Elsevier was done, with a focus on

Resident Poster # 162 Category: Clinical Vignette

Residency Program: Wayne State University Detroit Medical Center Sinai Grace
Presenter: Suneel Kumar
Additional Authors: Kausar Hafeez, MD; Zainab Muslehuddin, MD.

When chronic pancreatitis reaches the chest, a case of pancreaticopleural fistula and recurrent pleural effusion.

Introduction:

Pancreaticopleural fistula (PPF) is an infrequent complication arising in approximately 0.4% of pancreatitis cases, predominantly observed in patients with alcoholic chronic pancreatitis. This rare condition is characterized by an abnormal connection between the pancreas and the adjacent pleural cavity. It occurs due to a disruption of the pancreatic duct (PD), resulting in leakage of pancreatic fluid. Its symptoms primarily affect the thoracic region, posing challenges in diagnosis. Here, we describe a case of PPF in a

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 163 Category: Continuous Quality Improvement/Evidence-based Medicine

Residency Program: Wayne State University Detroit Medical Center Sinai Grace
Presenter: Anh Tuan Mai
Additional Authors: Malitha Hettiarachchi MD, FACP

Fixed versus Automatic CPAP in Adult Obstructive Sleep Apnea Management

Introduction:

Obstructive sleep apnea (OSA) involves repetitive pharyngeal airway collapse during sleep, leading to hypoxemia and hypercapnia. It is managed with either automatic positive airway pressure (APAP) or continuous positive airway pressure (CPAP). The objective of this study is to compare the efficacy of APAP and CPAP in the treatment of adult patients with OSA.

Methods:

A systematic review of literature from 2013-2023 in PubMed and Web of Science databases was conducted, focusing on APAP and CPAP in OSA management. Selection criteria included clinical trials, randomized controlled trials, and systematic reviews in patients greater than 18 years old. Duplicates were excluded via Rayyan.ai. Thematic analysis covered aspects such as pressure titration, symptom improvement, patient adherence, and adverse effects.

Results:

Seventeen studies met the inclusion criteria. In three studies, APAP was found to require significantly lower pressure settings for optimal titration. Both APAP and CPAP showed equivalent symptom improvement in all studies included. A slight hemodynamic advantage was seen with CPAP concerning diastolic in two studies and systolic pressures in one study. Higher BMI and male gender patients showed increased airway pressures with APAP. Patients with lighter sleep or higher arousal thresholds may tolerate and adhere to CPAP better. No notable differences in adverse effects between the two methods were identified.

Concl AP sp] TJ 5m50 0.0000912 0 612 792 re W* n BT /F2 10 Tf 1 0 0 1 181.08 539.67 Tm 0 g 0 G [( )] TJ ET Q

2024 Michigan Chapter ACP and SHM Michigan Chapter

Resident Poster # 166 Category: Clinical Vignette

Residency Program: Wayne State University Detroit Medical Center Sinai Grace
Presenter: Rakesh Shah, MD
Additional Authors: Priyanka Bhagat, MD; Khaled Harmouch, MD; Sumit Gami, MD; Kauser Hafeez, MD, MPH; Ahmed J Chaudhary, MD, FACP

Shifting paradigms of platinum-resistant ovarian cancer - mirvetuximab and bevacizumab, an elixir or curse? A case report of treatment-induced pulmonary fibrosis in a 68-year-old female with epithelial ovarian carcinoma.

Introduction

Mirvetuximab (MIRV) is the first biomarker-driven antibody-drug conjugate to target the folate receptor alpha (FRa) antigen - a protein expressed in 90% of ovarian malignancies. Clinical trials have demonstrated this drug to have significant efficacy in the treatment of platinum-resistant ovarian cancer (PROC). Despite the promising efficacy and tolerable safety profile, the drug when used with bevacizumab is associated with a rare and devastating treatment-related adverse effect of pulmonary fibrosis.

Case Description

A 68-year-old female presented to the emergency department for a 4-day duration of acute onset dyspnea with cough. She denied fever, hemoptysis, chest pain, or diaphoresis. Medical history was significant for high-grade serous ovarian carcinoma for which the patient had undergone neoadjuvant chemotherapy, cytoreductive surgery, adjuvant chemotherapy with 10 cycles of Bevacizumab, and maintenance therapy with Letrozole. She was later

Resident Poster # 167 Category: Continuous Quality Improvement/Evidence-based Medicine

Residency Program: Wayne State University Detroit Medical Center Sinai Grace
Presenter: Taher Shawartamimi
Additional Authors: Hettiarachchi, M

An Analysis of Ketamine Use in Mechanically Ventilated Patients: A Systematic Review

Introduction: Hemodynamic instability is one of the major side effects of many regularly used sedatives, which makes them problematic in the critical care situation. Ketamine is a new option that is gaining popularity due to its sedative effects on patients on mechanical ventilation. Nonetheless, a thorough analysis of its safety and adverse impact profile is necessary.

Materials and Methods: We searched the literature using PubMed with the terms "Ketamine" and "mechanical ventilation." The first search produced 34 results, and among the eligible trials were adult patients on mechanical ventilation who were continuously receiving an intraven.47 Tm ET Q -26(t)10(hods)-6(:)-6(W)-8(e s)-9(earche)-2(d t)11(he

Resident Poster # 168 Category: Clinical Vignette

Residency Program: Wayne State University Detroit Medical Center Sinai Grace
Presenter: Sujan Shrestha
Additional Authors: Amna Nadeem Kiani, MD; Naisargee Solanki, MD; Zainab Muslehuddin, MD; Ahmed Jamal Chaudhary, MD, FACP.

Uncommon Positional Dyspnea Manifestation Associated with Incidentally Diagnosed Adenomatous Goiter

Introduction:

Adenomatous goiter is a common benign tumor of the thyroid gland. Most cases are asymptomatic but various

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 169 Category: Clinical Vignette

Residency Program: Wayne State University Detroit Medical Center Sinai Grace
Presenter: Heitor Smiljanic Carrijo
Additional Authors: Ahmad Alnasarat, Juliana Chaves de Oliveira Paiva, Fahad Hanna.

Salmonella causing prostatic abscess; Rare case report

Introduction:

Prostatic abscesses (PA) are a rare urological condition associated with chronic or acute bacterial prostatitis[1]. Gram-negative bacilli (primarily E. coli) are the most prevalent, accounting for 50-60% of cases. Mostly, it is a diagnostic challenge as the patients present with non-specific symptoms, such as pelvic/perineal pain, dysuria, fever, chills, and urethral discharge. However, in this rare case, the cultured specimens revealed the presence of Salmonella Agona as the etiological agent. Salmonella is a rarely causative organism of prostatic abscess and is associated with an immunosuppressive state; in this case, the patient had uncontrolled Diabetes Mellitus type II.

We are reporting this case because the identification of a rare organism is vital: It changes the threshold for investigating complications common with this specific organism, such as enteric infections and bacteremia.

Case Description:

A 41-year-old male with a past medical history of uncontrolled T2DM s/p Bilateral below-the-knee amputation secondary to gangrenous diabetic foot infection, CKD, anxiety disorder, major depressive disorder, anemia, and chronic debility presented to the emergency department with complaints of pelvic and lower abdominal pain aggravated by recent constipation, pneumaturia, chills, and fever. The patient was hemodynamically stable during the initial examination. The patient had suprapubic pain on palpation and costovertebral angle tenderness on physical exam. Laboratory data did not reveal leukocytosis as expected. CT scan of the abdominal and pelvic revealed a large prostate abscess measuring 3.9 x 5.7 x 5.2 cm with bladder wall thickening and intraluminal air consistent with cystitis. Empirical antibiotic therapy was promptly initiated (Ceftriaxone on ED, then empirically changed to Vancomycin + Meropenem after CT report) to address this condition, and the patient underwent surgical intervention (transurethral unroofing) for abscess drainage and irrigation. Later, cultured specimens sent during the surgical procedure grew pure culture of Salmonella Agona. He also received adequate control of his glucose level. Over the following week, the patient's condition improved; EGD for possible GI bleed, as the patient mentioned a melena episode, showed mild esophagitis-gastritis-duodenitis, which may be associated with the nontyphoidal Salmonella as it is known to cause enteric infection, the patient was started on Pantoprazole 40 mg daily with the improvement of his symptoms. After 7 days as an inpatient, he was discharged home with a

g 0 G [(chhi)6(s)-ha5eontywt] TJ tti. Aftta17 + Mse enteric infecnteric infecnteric infec consi. n 0 G [(chhi)6(s692 re W* n

Resident Poster # 170 Category: Research

Residency Program: Wayne State University Detroit Medical Center Sinai Grace
Presenter: Nishita Tripathi
Additional Authors: Dmitry Chernyshev, MD, Shifaatullah K.S. Mohammed, MD, Fred H.D. Charles, MD, MPH, Malitha Hettiarachchi, MD, FACP

Solitary plasmacytoma of bone presenting as atypical chest pain in an elderly male: A case report

Introduction: Plasma cell dyscrasias comprises a broad spectrum of disorders characterized by neoplastic proliferation of clonal plasma cells. Solitary plasmacytoma of bone (SPB) contains a single clone of plasma cells localized to a single bone lesion witale: A case report

2024

Resident Poster # 172 Category: Clinical Vignette

Residency Program: Wayne State University Providence Rochester Hospital
Presenter: Nour Aldaoud
Additional Authors: Gagandeep Singh (MD), ET Q qBU( )] TJ Ed

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 175 Category: Continuous Quality Improvement/Evidence-based Medicine

Residency Program: Wayne State University Providence Rochester Hospital
Presenter: JAIKUMAR
Additional Authors: Ayushi Garg (MD), Jessica Lutkenhoff (MS3), Sarwan Kumar (MD)

"The Role of Residents in Patient Safety: Evaluating Rapid Response Outcomes in a Community-Based Teaching Hospital"

Introduction:

In response to the high incidence of cardiopulmonary arrests, a leading cause of inpatient mortality, hospitals have implemented rapid response systems. This retrospective cohort study examines the role of residents in patient safety and the outcomes of rapid responses in a community-based teaching hospital.

Method:

Data from May 2022 to October 2023 were analyzed to assess the impact of rapid responses on inpatient mortality, code blues, and unexpected transfers to the ICU. The study aimed to standardize the diagnosis and treatment of patients with SIRS 2/4 as sepsis, aiming to reduce rapid responses due to SIRS >2/4 by 25% within six months. The analysis focused or-3(sF2 10 Tf 1 0 0 1 193.58 515.15 Tm 0 g 0 G [(pi)5(d res)-10(pons)-6(es)-10(i)6(n a)5

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day

Resident Posters

Resident Poster # 176 Category: Continuous Quality Improvement/Evidence-based Medicine

Residency Program: Wayne State University Providence Rochester Hospital
Presenter: Vamsi Krishna Lavu
Additional Authors:

2024

2024

Resident Poster #

Resident Poster # 181 Category: Clinical Vignette

Residency Program: Western Michigan University Homer Stryker M.D. School of Medicine
Presenter: Ayema Haque
Additional Authors: Alec Johnson, Bruna Dellatore Diniz, Ali Shammout, Joanne Baker

Group C Streptococcus Species: A rare cause of Infective endocarditis

Infective endocarditis is defined by infection of the endocardial surface of the heart, heart valves, or indwelling

2024 Michigan Chapter ACP and SHM Michigan Chapter Resident and Medical Student Day
Resident Posters

Resident Poster # 183 Category: Clinical Vignette

Residency Program: Western Michigan University Homer Stryker M.D. School of Medicine
Presenter: Alana Pinheiro Alves
Additional Authors: Alana Pinheiro Alves MD, Susan Bannon, MD, Ibrahim Zahid

Sporadic Creutzfeldt-Jakob Disease with Visual Disturbance: A Rare Presenting Symptom

Background:

Creutzfeldt-Jakob disease (CJD) is a rapidly advancing neurodegenerative condition marked by dementia, myoclonus, and other neurological manifestations. While visual symptoms are present in 20% of CJD cases, very few patients present with an isolated visual disturbance without noticeable cognitive decline initially; we present one such case in our report.

Case report:

A 60-year-old woman presented to the emergency department (ED) for visual disturbance, worsening confusion, abnormal gait, and urinary retention. She had a history of chronic left visual impairment. Symptoms of decreased vision in the right eye had started about two months before: she had seen an ophthalmologist about four weeks after symptoms began, and a slit lamp and tonometry tests were normal; given normal C-reactive protein, serum copper, and B12 vitamin levels, she was advised to monitor symptoms at home. On three weeks follow-up, she had decreased visual acuity of 20/40 in her right eye, following which she developed progressive difficulty with word finding, memory, and wobbly gait, as well as urinary retention. In the ED, her physical exam was significant for lack of coordination, confusion, diffuse bradykinesia, abnormal extraocular eye movements, and hyperreflexia of left lower extremity, with significant motion tremors on finger-to-nose testing and diffuse spasticity. Basic blood work and computed tomography imaging of brain were largely unremarkable. MRI of the brain with and without contrast however showed mildly scattered white matter hyperintensities. Shortly after admission, the patient became non-verbal and had multiple episodes of repeated tonic-clonic seizures requiring intubation. An electroencephalogram (EEG) was performed, which showed several instances of sharp and slow waves suspicious of electrographic seizure. In the ICU post-intubation, a CSF analysis was performed, remarkable for RBCs 6831 per u/L, with 10 WBCs per u/L (100% mononuclear), glucose 65 mg/dl, and protein 40 mg/dl, with negative culture, VDRL, Lyme PCR, oligoclonal bands, paraneoplastc antibodies and malignant cells. 14-3-3 protein testing on the CSF was initially inconclusive, but tau-protein was high (>20000 pg/ml). The initial Real-time quaking-induced conversion (RT-QuIC) assay was negative. Unfortunately, her encephalopathy progressed; repeat MRI of the brain with and without contrast showed cortical ribboning. Given the lack of diagnostic answers, repeat CSF collection was performed, and the RT-QuIC test resulted positive, confirming CJD. With family input she was transitioned to comfort care measures, extubated and passed away.

Discussion:

CJD is a disorder of poor prognosis and may be acquired through medical procedures, exposure to highly contaminated tissue or inherited genetically, but ~80% cases are categorized as sporadic CJD, where the underlying cause remains unknown as in our case. The most common clinical manifestations of CJD are progressive cognitive decrease, dementia, myoclonus, and ataxia. When the disease involves predominant visual symptoms, it is termed as the “Heidenhain variant”. It must be considered a differential diag